

- Ficarro, S.B., McClelland, M.L., Stukenberg, P.T., Burke, D.J., Ross, M.M., Shabanowitz, J., Hunt, D.F. and White, F.M. *Nature Biotechnology*, 2002, 20, 301-305.
- Gruhler A, Olsen JV, Mohammed S, Mortensen P, Faergeman NJ, Mann M, Jensen ON. *Mol Cell Proteomics*. 2005 Mar;4:310-27. Epub 2005 Jan 22.
- Li X, Gerber SA, Rudner AD, Beausoleil SA, Haas W, Villén J, Elias JE, Gygi SP. *J Proteome Res*. 2007 Mar; 6:1190-7
- Connor P.A. & Dobson K.D. *Interface. J Langmuir* 1999, 15, 2402-2408.
- Larsen, M.R., Thingholm, T.E., Jensen, O.N., Roepstorff, P. & Jorgensen, T.J. *Mol Cell Proteomics* 2005, 4, 873-886.
- Zhang X, Ye J, Jensen ON, Roepstorff P. *Mol Cell Proteomics*. 2007 Nov;6(11):2032-42. Epub 2007 Aug 4.
- Thingholm TE, Jensen ON, Robinson PJ, Larsen MR. SIMAC . *Mol Cell Proteomics*. 2007 Nov 26.
- Hjernø K. Chapter of Ph.D. Thesis by Karin Hjernø. Protein Research Group, Department of Biochemistry and Molecular Biology, University of Southern Denmark, Odense.
- Mann M, Ong SE, Grønborg M, Steen H, Jensen ON, Pandey A. *Trends Biotechnol*. 2002 Jun;20(6):261-8.
- Jensen ON. *Nat Rev Mol Cell Biol*. 2006 Jun;7(6):391-403.
- Ingrell CR, Miller ML, Jensen ON, Blom N. *Bioinformatics*. 2007 Apr 1;23(7):895-7. Epub 2007 Feb 5.
- Schulze WX, Mann M. *J Biol Chem*. 2004 Mar 12;279(11):10756-64. Epub 2003 Dec 16.
- Matthiesen R. *Virtual Expert. Methods Mol Biol*. 2007;367:121-38.

## New insights in the study of s-nitrosylation & s-nitration: strategies and problems

Martínez-Acedo P<sup>1</sup>, Martínez-Ruiz A<sup>2</sup>, Maldonado AM<sup>3</sup>, Horcajo-Redondo M<sup>4</sup>, Jorge I<sup>1</sup>, Serrano I<sup>1,5</sup>, Navarro PJ<sup>1</sup>, Pérez-Hernández D<sup>1</sup>, Nuñez E<sup>1</sup>, Redondo JM<sup>2</sup>, Jorrín JJ<sup>3</sup>, Lamas S<sup>4</sup>, Vazquez J<sup>1</sup>.

<sup>1</sup>Protein Chemistry and Proteomics Laboratory, Centro de Biología Molecular Severo Ochoa, CSIC-UAM, Madrid, Spain, <sup>2</sup> Centro Nacional de Investigaciones Cardiovasculares, CNIC, Madrid, Spain, <sup>3</sup> Universidad de Córdoba, Córdoba, Spain, <sup>4</sup> Centro de Investigaciones Biológicas, CNIC, Madrid, Spain, <sup>5</sup> Biology Department, University of Puerto Rico, Arecibo Campus, Puerto Rico, USA.

### Introduction

The study of the physiological role of reactive nitrogen species (RNS) is attracting increasing interest in the biomedical community. The most common protein modifications by RNS are S-nitrosylation or covalent incorporation of a nitrosile radical to a thiol group, and nitration, or incorporation of a NO<sub>2</sub> group, generally to tyrosine residues. It is currently thought that nitrosative stress is one of the most relevant pathogenic factors in cardiovascular pathology, including atherosclerosis, hypertension, diabetes and cardiac insufficiency, and NO has be-

come recognized as a key signaling molecule in plants. Moreover, there are evidences that exposing endothelial cells to the immunosuppressor cyclosporin A (CsA) increases production of RNS, and particularly peroxynitrite, that stimulates nitration of proteins, being the protein MnSOD one of the candidates to CsA-induced nitration (Horcajo-Redondo *et al.*, 2005). Here we have developed and used proteomics techniques to identify specific sites of S-nitrosylation and S-nitration in protein extracts from different species. The growing interest in the study of these modifications have led to the creation of a Spanish working group, the "Nitrosoteam",

with the objective of increasing the knowledge of the modifications in different kinds of samples by using a common strategy.

**Material and methods**

We used the “biotin-switch” method (Martínez-Ruiz *et al.*, 2005) to selectively enrich nitrosylated peptides in human and plant protein extracts. The Selected MS/MS Ion Monitoring (SMIM) technique using a linear ion trap mass spectrometer (LTQ, Thermo-Fischer Scientific, San Jose, USA) (Jorge *et al.*, 2007) was used to identify a specific site of nitration in a MnSOD peptide.

**Results**

Here we present the first results of the “Nitroso-team”. The first goal was to apply the “biotin-switch” method to the purification of peptides instead of proteins, what is an innovation of the technique. We first used synthetic peptides as probes and after optimization of the mass spectrometer and the database searching conditions we proceeded to the study of *in vitro* cell culture extracts without blocking free cysteine thiols with MMTS. Elution of peptides from the avidin column was made with either acid or reducing conditions. The acidic elution had the advantage that biotin remained attached to the peptide, whereas in the reducing elution more peptides could be identified. Some problems in the different steps of the experiment have emerged. We observed that the biotin moiety attached to the peptides produced a lot of fragments that could not be assigned (Figure 1). Besides, the manipulation of the samples produced different atypical adducts that make the identification more difficult. For studying the nitration of a specific site in MnSOD, we had to optimize the mass spectrometer conditions in order to identify beyond any doubt the modification site using multiple fragmentation stages (MS<sup>3</sup>).


**Conclusions**

Several technical problems were resolved during the course of this study to analyze S-nitrosyla-

tion and S-nitration sites by MS approaches. Our preliminary results give us good indications about the viability and specificity of the technique and encouraged the “Nitroso-team” to apply these developments to the analysis of physiological samples.

**References**

Jorge I., Casas E.M., Villar M., Ortega-Pérez I., López-Ferrer D., Martínez-Ruiz A., Carrera M., Marina A., Martínez P., Serrano H., Cañas B., Were F., Gallardo J.M., Lamas S., Redondo J.M., García-Dorado D., Vázquez J. *J Mass Spectrom.* 2007, 42(11): 1391-1403.  
 Martínez-Ruiz A., Lamas S. *Methods Enzymol.* 2005, 396: 131-139. Redondo-Horcajo M., Lamas S. *J Nephrol.* 2005, 18(4): 453-457.



**Figure 1:** Fragmentation spectra of a non biotinilated (upper panel) and a biotinilated peptide (lower panel) showing the presence of multiple fragments produced by biotin that could not be assigned (circles).