

## Article

# Short and Long-Term Effect of Land Use and Management on Soil Organic Carbon Stock in Semi-Desert Areas of North Africa-Tunisia

Fatma Baraket <sup>1</sup>, Manuel González-Rosado <sup>2</sup> , Nadhem Brahim <sup>3</sup> , Núria Roca <sup>4</sup> , Hadda Ben Mbarek <sup>1</sup>, Marcin Switoniak <sup>5</sup> , Rayda Chaker <sup>6</sup>, Ángel Sánchez Bellón <sup>7</sup> , Hafedh Rigane <sup>1</sup>, Kamel Gargouri <sup>6</sup>  and Luis Parras-Alcántara <sup>2,\*</sup> 

- <sup>1</sup> Research Laboratory of Environmental Sciences and Sustainable Development (L18ES32), Earth Sciences Department, Sfax Faculty of Sciences, P.O. Box 1087, Sfax 3000, Tunisia; baraketfatouma@gmail.com (F.B.); haddabenmbarek@gmail.com (H.B.M.); hafedhrigane@yahoo.fr (H.R.)
- <sup>2</sup> SUMAS Research Group, Department of Agricultural Chemistry, Soil Science and Microbiology, Faculty of Science, Agrifood Campus of International Excellence—ceiA3, University of Cordoba, 14071 Cordoba, Spain; mgrosado@uco.es
- <sup>3</sup> Department of Geology, Faculty of Sciences of Tunis, University of Tunis El Manar, Tunis 2092, Tunisia; nadhem.brahim@fst.utm.tn
- <sup>4</sup> Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona, Av. Diagonal 643, 08028 Barcelona, Spain; nroca@ub.edu
- <sup>5</sup> Department of Soil Science and Landscape Management, Nicolaus Copernicus University in Toruń, 1 Lwowska Str., 87-100 Toruń, Poland; swit@umk.pl
- <sup>6</sup> Olive Institute, University of Sfax, P.O. Box 1087, Sfax 3000, Tunisia; raydachaker@gmail.com (R.C.); kamel.gargouri@iresa.agrinet.tn (K.G.)
- <sup>7</sup> Department of Earth Sciences, Faculty of Sciences, Campus University of Puerto Real, Cadiz University, 11519 Puerto Real, Spain; angel.sanchez@uca.es
- \* Correspondence: luis.parras@uco.es; Tel./Fax: +34-9-5721-1092



**Citation:** Baraket, F.; González-Rosado, M.; Brahim, N.; Roca, N.; Mbarek, H.B.; Switoniak, M.; Chaker, R.; Bellón, Á.S.; Rigane, H.; Gargouri, K.; et al. Short and Long-Term Effect of Land Use and Management on Soil Organic Carbon Stock in Semi-Desert Areas of North Africa-Tunisia. *Agriculture* **2021**, *11*, 1267. <https://doi.org/10.3390/agriculture11121267>

Academic Editor: Jacopo Bacenetti

Received: 30 September 2021  
Accepted: 5 December 2021  
Published: 14 December 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Soil organic carbon (SOC) plays an important role in the global C cycle, as well as in the maintenance and improvement of the soil quality. Over time, special attention has been paid to it in the study of the SOC reserves worldwide; however, reduced attention has been given to assessing the spatial patterns of SOC stock (SOCS) in semi-desert ecosystems. In this line, there are no conclusive studies in drylands of Africa affected by aeolian processes (semi-desert conditions) mainly due to the complexity of sample collection, and this is especially significant in some soil types such as Arenosols (AR) and Calcisols (CL). This study evaluated the spatial variability of SOC and SOCS in AR and CL with woody crops in relation to land use and management (old plantations > 100 years: centenary olive grove; new plantations < 12 years: young olive grove, almond, and pistachio) in semi-desert conditions. For this purpose, 16 soil profiles (for 0–40 and 40–100 cm depth) were selected and studied in an experimental area of Menzel Chaker-Sfax in southeastern Tunisia (North Africa). The main results indicated that the SOCS on average was higher in Old Cultivated AR (OC-AR) with 41.16 Mg ha<sup>-1</sup> compared to Newly Cultivated AR (NC-AR) with 25.13 Mg ha<sup>-1</sup>. However, the SOCS decreased after a long period of cultivation in CL from 43.00 Mg ha<sup>-1</sup> (Newly Cultivated CL: NC-CL) to 32.19 Mg ha<sup>-1</sup> (Old Cultivated CL: OC-CL). This indicates that in the long term, CL has more capacity to store SOC than AR, and that in the short term, AR is more sensitive to land management than CL.

**Keywords:** arid climate; land use; land management; Calcisols; Arenosols; soil organic carbon stocks; olive grove; almond grove; pistachio grove

## 1. Introduction

In most terrestrial ecosystems, soil organic carbon (SOC) is the largest carbon (C) pool [1] containing approximately 2344 Gt of organic C globally [2]. In fact, SOC is an

essential element of earthly life and constitutes the most fertilizing element of the soil, and SOC content is an important indicator of biological and microbiological activity [3,4]. Therefore, SOC and soil organic matter (SOM) are considered major factors that influence the soil quality, affecting agricultural productivity, soil structural stabilization, plant nutrients retention, and water-holding capacity, among other things [5,6]. However, in addition, SOC stock (SOCS) depends on factors such as wildlife, earthworms, ants, and microorganisms such as fungi and bacteria and organic waste [7], and this SOCS can be found at 0–2 m depth [8,9]. Other factors that can affect to SOCS are the climatic conditions (precipitation and temperature) and agriculture practices (land use and management) [10]. In this sense, high temperatures, low rainfall, and intensified tillage can accelerate SOM decomposition, conditioning agricultural production in many regions of the world [11]. Therefore, a decrease in the SOM content leads to reduction in soil fertility and productivity [12]. Therefore, SOCS should be estimated to evaluate the impact of land management practices on soil fertility and in the greenhouse gases generation [13,14].

All these considerations are especially important in drylands (semi-arid, arid, semi-desert, and desert areas) with low average annual rainfall (<500 mm) [15], with a cover more than one-third of the Earth's land surface, and with more than 36% of the world's SOCS [16,17]. In this line, [18] indicates that these ecosystems can be more responsive to elevated CO<sub>2</sub> than others because net primary productivity is mostly limited by water availability. Additionally, other authors such as [19] have reported that desert soils exhibit higher atmospheric CO<sub>2</sub> fixation capacity than other soils (e.g., meadow soils).

In this sense, researchers in [20] indicated that dryland areas in Tunisia is characterized by low SOCS (18.7 Mg ha<sup>-1</sup>) due to sandy texture, which is associated with low SOM content and reduced cation exchange capacity [21,22], highlighting that in sandy soils, the SOM decomposition has a higher rate than in soils with high clay content [23]. However, water scarcity is also considered as another crucial factor that causes soil degradation, and a decrease in SOC content in semi-arid and arid regions [24]. On average, every year, arid areas have long drought periods (4 to 6 months), so the generation of natural biomass as C source is limited by climatic conditions, but this problem is maximized due to management practices in these arid regions, since the highly intensified tillage causes a loss of spontaneous vegetation cover [25]. The main consequence of these processes acting synergistically is that these soils are highly vulnerable to degradation processes, due mainly to SOM depletion and wind erosion, which cause desertification, which is intensified by the current global warming crisis [26,27].

Recently, many researchers have shown great interest in the SOCS prediction (at regional scale), extrapolating the SOCS on a wide range of soil and/or vegetation types in arid areas of Africa [20]. In this line, [28] analyzed the SOC storage evolution in relation with tillage system in three soil types (Vertisols, Cambisols, and Luvisols) under Mediterranean climate, emphasizing that the SOC was significantly higher in no tillage compared to conventional tillage (10% more in Vertisols and 8% more in Cambisols), with no significant differences in the Luvisols case. More recently, [29] studied the SOCS distribution in Cambisols under different crops in semi-arid Tunisian climate, suggesting that the SOCS has exceeded the national and international standards in Cambisols. Additionally, in Tunisia, SOCS was assessed in different soil types including Lithosols, Solonchaks, Cambisols, and Regosols [20,29,30]; however, Calcisols (CL) and Arenosols (AR) have not been studied yet. In this line, is important to note that three quarters of Tunisia is under arid or semi-desert climate. Therefore, analyzing the SOCS in CL and AR in Tunisian is important to improve the soil quality and its productive capacity, in addition to being able to extrapolate these data to other semi-desert areas to intervene in the land use and management of these soil types. In this sense, several studies have established that soils in arid and semi-desert regions after cultivation suffer significant variations in their biological, physical, and chemical properties [31–35]. However, little research has focused on assessing the effects of woody crops (olive, almond, and pistachio) at different development stages in AR and CL during short- and long-term periods and their relationship with the soil properties and


SOCS. Therefore, due to the role that these soils play in C sequestration and in the crop productivity, it is essential to understand the effectiveness of land use and managements under extreme climatic conditions not only in terms of physical properties but also in the C dynamic variation.

Given this scenario, the general objectives of this study are (i) to study the SOC in Calcisols and Arenosols of Tunisia (dryland rainfed area) and (ii) to analyze in the short and long terms the soil's capacity for SOC stock under different land uses and management.

## 2. Materials and Methods

### 2.1. Study Area

The study area is in an experimental farm of Essalema, located at 50 km in the north-west in Sfax governorship, in the Menzel Chaker delegation—Tunisia ( $34^{\circ}59'15''$  N– $10^{\circ}20'03''$  W) (Figure 1). This area covers a surface of 18,670 ha and is divided into seven sub-farms, with an average altitude of 161 m.a.s.l., ranging from 105 m.a.s.l. to 217 m.a.s.l. (meters above sea level), with slopes < 3% slightly undulating.



**Figure 1.** Maps of the study area: (a) Tunisia map, (b) Sfax location, and (c) Menzel-Chaker district with the location of soil profiles <https://earth.google.com/web/@36.6638314,7.9704552,553.33197981a,3949044.97759938d,35y,0h,0t,0r> (Google Earth Pro—Access: 27 September 2021).

The climate is arid Mediterranean and hot arid steppe according to the Köppen–Geiger updated classification [36]. The annual average temperature was  $19.5^{\circ}\text{C}$ , with a maximum air temperature of  $33^{\circ}\text{C}$  in August and a minimum air temperature of  $6^{\circ}\text{C}$  in January (temperature data provided by the National Institute of Metrology for the period 1966–2019). The annual average precipitation was 169 mm and monthly rainfall ranges

from 6 mm (July) to 34 mm (October) (rainfall data provided by Salama farm station for the period 2008–2019).

The study area is characterized by outcrops of the Middle and Upper Continental Pleistocene, and the Lower Pleistocene, formed by limnic sabkhas (deposits of coastal flats subject to periodic flooding and evaporation) and recent alluvium. Most of the wadis are endorheic, leading to closed depressions of the Sebkhass and Garâas type. Depending on their morpho-structural conditions, these closed depressions take the form of synclinal basins (Menzel Chaker region) or the form of Sebkhass sand Garâas (Bou Jmal, Karafita). In the dryland regions, the aeolian processes also play an important role, particularly where the precipitation is low (<150 mm y<sup>-1</sup>) [37,38], so the aeolian materials accumulation leads to the poorly developed sandy soils formation [39], as it happens in the study area's experimental field [40].

## 2.2. Soil Sampling and Analytical Methods

Samples from 16 soil profiles were collected: 10 in CL and 6 in AR (these soil types covered most of the studied area ~70%). All investigated soils were tilled to depths varying between 20 and 30 cm (can reach up to 40 cm deep) four to six times per year using a tractor with cultivator depending on rainfall. Soil profiles were dug with a mini excavator, and soil samples were collected at different soil control section for each soil profile (S1: 0–40 cm; S2: 40–100 cm), for a proper determination of physical and chemical soil properties [41,42].

The collected samples were labeled (A1–A5: samples of the Newly Cultivated Calcisols (tilled < 12 years)—NC-CL; M1–M5: samples of the Old Cultivated Calcisols (tilled > 100 years)—OC-CL; C1–C3: samples of the Newly Cultivated Arenosols (tilled < 12 years)—NC-AR; B1–B3: samples of the Old Cultivated Arenosols (tilled > 100 years)—OC-AR).

Soil samples were placed in polyethylene bags, which were labeled and transferred to the laboratory and air dried. Once dried, the samples were sieved at 2000 µm, separating the thick fragments and roots from the rest of the material. Three repetitions were carried out for each sample. The analytical methods, laboratory analysis, and other parameters calculated used in this study to determine different soil properties are reported in Table 1, according to handbook of plant and soil analysis for agricultural systems [43]. Soils were described and classified according to World Reference Base for Soil Resources [38].

**Table 1.** Analytical methods used in this study (field measurements, laboratory analysis, and parameters calculated).

Parameters	Method
Field measurements	
Bulk density (Mg m <sup>-3</sup> )	Core method [44] <sup>a</sup>
Laboratory analysis	
Particle size distribution	Robinson pipette method [45] <sup>b</sup>
pH—H <sub>2</sub> O	Suspension in water 1:2.5 [46]
Total Organic C (g kg <sup>-1</sup> )	Walkley and Black method [47]
CaCO <sub>3</sub> (%)	Soil Calcium carbonate equivalent [48,49]
Parameters calculated	
SOC-S (Mg ha <sup>-1</sup> )	SOC-S = SOC concentration × BD × d × (1 - δ2 mm%) × 10 <sup>-1</sup> [50–52] <sup>c</sup>
T-SOC-S (Mg ha <sup>-1</sup> )	T-SOC-S = Σ <sub>soil horizon 1 ... n</sub> SOC-S <sub>soil horizons</sub> [52] <sup>d</sup>

For all the parameters studied, the recommendations of the Handbook of Plant and Soil Analysis for Agricultural Systems have been followed [43]. <sup>a</sup> 3 cm in diameter, 10 cm in length, and 70.65 cm<sup>3</sup> in volume. <sup>b</sup> Prior to determination of particle size distribution, samples were treated with H<sub>2</sub>O<sub>2</sub> (6%) to remove organic matter (OM). Particles larger than 2 mm were determined by wet sieving, and smaller particles were classified according to USDA standards (2004). <sup>c</sup> Where SOC is the organic carbon content (g kg<sup>-1</sup>), d is the thickness of the soil layer (cm), δ2mm is the fractional percentage (%) of soil mineral particles >2 mm in size in the soil, and BD is the soil bulk density (Mg m<sup>-3</sup>). <sup>d</sup> T-SOC-S: Total SOC stock determined by adding all the soil horizons considered.

## 2.3. Experimental Design

The research was carried out in an experimental farm in Essalema–Sfax–Menzel–Tunisia. For this, two soil types were selected (AR and CL), with different land uses (OG: olive grove,

AT: almond and PT: pistachio) and managements. The soils' evolution in the short (<12 years) and long terms (>100 years) was compared to analyze the effects of management on SOCS. Land use and management are characterized in Tables 2 and 3 and Figure 2.

**Table 2.** Land use, managements, and soil qualifiers of the soils (Arenosols) in the study area.

Soil	SP	Soil Type	Hor	Depth (cm)	Land Use	Management
AR	OC-AR	B1	Ap AC	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture; Olive pomace 3.5 Mg ha <sup>-1</sup> every 10 years
		B2	ARca C	0–40 40–100	OG	Density: 20 trees ha <sup>-1</sup> ; Amendments: Olive pomace 3.5 Mg ha <sup>-1</sup> every 10 years
		B3	AREu AC	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture; Olive pomace 3.5 Mg ha <sup>-1</sup> every 10 years
	NC-AR	C1	ARca C	0–40 40–100	AT	Density: 68 trees ha <sup>-1</sup> ; The plowing is Canadian type (plowing just with the tails of the washers). Without organic amendment
		C2	ARca BC	0–40 40–100	OG	Density: 34 trees ha <sup>-1</sup> ; Without organic amendment
		C3	AREu C	0–40 40–100	OG	Density: 34 trees ha <sup>-1</sup> ; Without organic amendment

ARca: Calcaric Arenosols; AREu: Eutric Arenosols; SP: Soil profile; OC-AR: Old cultivated (tilled > 100 years) Arenosols; NCAR: Newly cultivated (tilled < 12 years) Arenosols; SP: Soil Sampling; Hor: Horizon; OG: Olive grove; AT: Almond tree. C2 and C3: The plowing is Canadian type (plowing just with the tails of the washers when the OG is big, sometimes using the blade types. They are plowed 4 times/year.

**Table 3.** Land use, managements, and soil qualifiers of the soils (Calcisols) in the study area.

Soil	M	SP	Soil Type	Hor	Depth (cm)	Land Use	Management
CL	OC-CL	M1	ha CL	Ap 2Ck	0–40 40–100	OG	Density: 20 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture 3.5 Mg ha <sup>-1</sup> every 10 years
		M2	ha CL	Ap 2Bk	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture; Olive pomace 3.5 Mg ha <sup>-1</sup> every 10 years
		M3	ha CL	Ap Ck	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; No Amendments
		M4	ha CL	Apk 2Ck	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; No Amendments
		M5	ha CL	Ap 2Ck	0–40 40–100	OG	Density: 17 trees ha <sup>-1</sup> ; No Amendments
	NC-CL	A1	ha CL	Ap 2Ck	0–40 40–100	OG	Density: 34 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture 1 Mg ha <sup>-1</sup> every 10 years
		A2	ha CL	Ap Ck	0–40 40–100	OG	Density: 34 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture 1 Mg ha <sup>-1</sup> every 10 years
		A3	ha CL	Ap 2Ck	0–40 40–100	OG	Density: 34 trees ha <sup>-1</sup> ; Amendments: sheep manure mixture 1 Mg ha <sup>-1</sup> every 10 years
		A4	ha CL	Ap Ck	0–40 40–100	AT	Density: 68 trees ha <sup>-1</sup> ; The plowing is Canadian type (plowing just with the tails of the washers)
		A5	ha CL	Ap 2Bk	0–40 40–100	PT	Density: 39 trees ha <sup>-1</sup> ; The plowing is Canadian type (plowing just with the tails of the washers)

ha CL: Haplic Calcisols; OC-CL: Old cultivated (tilled > 100 years) Calcisols; M: Management; SP: Soil Profile; NC-CL: Newly cultivated (tilled < 12 years) Calcisols; SP: Soil Sampling; Hor: Horizon; OG: Olive grove; AT: Almond tree; PT: Pistachio tree. A1, A2, and A3: The plowing is Canadian type (plowing just with the tails of the washers when the OG will be big, they use sometimes the blade types. They are plowed 4 times/year. M1, M2, M3, M4, and M5: The plowing is of the Canadian type. In the fall, they use the tails of the two-row pucks, in the winter they use the blade (teeth), one row only, and in the spring, they use the tails of the pucks (two rows) for weeding. During the summer, they use the blade (teeth), a single row.



**Figure 2.** Different land uses in the experimental farm: Centenary olive grove, young olive grove, almond tree, and pistachio tree. (1): Centenary olive grove—old cultivated > 100 years; (2): young olive grove—newly cultivated < 12 years; (3): almond tree—newly cultivated < 12 years; (4): pistachio tree—newly cultivated < 12 years.

All soil samples were taken at the same time (synchronic approach) under different management practices at known durations from an initial reference state, and the SOCS was compared under this initial reference state [53].

#### 2.4. Statistical Analyses

The effect of land management and soil depth on soil properties was analyzed using SPSS 20.0 for Windows. Data were tested for normality to verify the model assumptions using Duncan's multiple range tests, and differences of  $p < 0.05$  were considered statistically significant.

### 3. Results and Discussion

#### 3.1. Soil Characterization

The soils studied are developed from loamy or sandy material often of aeolian origin, and all soils had simple morphology with two genetic horizons—Ap horizon and underlying horizons with calcium carbonates accumulation (Ck and Bk); according to [38], the studied soils were mainly Haplic CL (CLha), Calcaric AR (ARca) and Eutric AR (AReu) (Tables 2–5).

**Table 4.** Principal soil properties evaluated (average  $\pm$  SD \*) in the soil profile by soil control section in Arenosols.

Soil	M	SP	LU	Hor	Depth (cm)	Gravel (%)	Sand (%)	Silt (%)	Clay (%)	BD (Mg m <sup>-3</sup> )	pH (H <sub>2</sub> O)	CaCO <sub>3</sub> (%)	OC (%)
AR	OC-AR <i>n</i> = (3 × 2)	B1	OG	Ap	0–40	1.15	96.37	1.17	2.46	1.701	9.18	2.64	0.125
				AC	40–100	0.60	92.99	4.62	2.39	1.645	9.14	2.78	0.330
		B2	OG	Ap	0–40	1.91	94.66	1.69	3.65	1.692	9.25	4.06	0.290
				C	40–100	6.74	92.95	2.44	4.61	1.673	9.19	7.42	0.171
		B3	OG	Ap	0–40	1.48	98.22	0.34	1.44	1.705	9.09	0.51	0.282
				AC	40–100	2.60	98.05	0.49	1.46	1.681	9.10	0.55	0.290
	X		Ap	0–40	1.51 $\pm$ 0.28	96.42 $\pm$ 1.78	1.07 $\pm$ 0.68	2.52 $\pm$ 1.11	1.70 $\pm$ 0.01	9.17 $\pm$ 0.08	2.40 $\pm$ 1.79	0.23 $\pm$ 0.09	
			AC	40–100	3.31 $\pm$ 3.13	94.66 $\pm$ 2.93	2.52 $\pm$ 2.07	2.82 $\pm$ 1.62	1.67 $\pm$ 0.02	9.14 $\pm$ 0.05	3.58 $\pm$ 3.51	0.26 $\pm$ 0.08	
	NC-AR <i>n</i> = (3 × 2)	C1	AT	Ap	0–40	2.98	80.51	10.75	8.74	1.560	8.95	4.11	0.270
				C	40–100	4.32	82.58	8.38	9.04	1.530	8.94	6.49	0.165
		C2	OG	Ap	0–40	0.66	88.91	5.22	5.87	1.632	8.81	3.03	0.290
				BC	40–100	0.88	95.32	1.33	3.35	1.681	8.68	2.63	0.095
C3		OG	Ap	0–40	0.09	94.01	2.23	3.76	1.670	9.12	1.04	0.145	
			C	40–100	0.44	91.58	2.09	6.33	1.690	9.17	0.81	0.065	
X		Ap	0–40	1.24 $\pm$ 1.53	87.81 $\pm$ 6.82	6.07 $\pm$ 4.32	6.12 $\pm$ 2.50	1.62 $\pm$ 0.06	8.96 $\pm$ 0.16	2.73 $\pm$ 1.56	0.24 $\pm$ 0.08		
		C	40–100	1.88 $\pm$ 2.13	89.83 $\pm$ 6.55	3.93 $\pm$ 3.87	6.24 $\pm$ 2.85	1.63 $\pm$ 0.09	8.93 $\pm$ 0.25	3.31 $\pm$ 2.90	0.11 $\pm$ 0.05		

SD \*: Standard deviation; M: Management; SP: Soil sampling; Hor: Horizon; BD: Bulk density; OC: Organic carbon; *n* = Sample size. OC-AR: Old cultivated (tilled > 100 years) Arenosols; NC-AR: Newly cultivated (tilled < 12 years) Arenosols.

**Table 5.** Principal soil properties evaluated (average  $\pm$  SD \*) in the soil profile by soil control section in Calcisol.

Soil	M	SP	LU	Hor	Depth (cm)	Gravel (%)	Sand (%)	Silt (%)	Clay (%)	BD (Mg m <sup>-3</sup> )	pH (H <sub>2</sub> O)	CaCO <sub>3</sub> (%)	OC (%)
CL	OC-CL <i>n</i> = (3 × 2)	M1	OG	Ap	0–40	6.91	91.43	3.77	4.80	1.621	9.26	5.66	0.131
				2Ck	40–100	7.47	74.28	8.69	17.03	1.591	9.10	20.47	0.201
		M2	OG	Ap	0–40	0.81	92.14	2.85	5.01	1.610	9.19	4.27	0.260
				2Bk	40–100	2.38	56.41	21.17	22.42	1.342	8.81	22.03	0.295
		M3	OG	Ap	0–40	23.54	73.98	12.89	13.13	1.330	8.78	11.61	0.540
				Ck	40–100	41.86	63.59	14.45	21.96	1.451	8.99	21.97	0.230
	M4	OG	Apk	0–40	16.16	90.97	5.79	3.24	1.570	9.12	10.08	0.115	
			2Ck	40–100	6.20	63.32	3.94	32.74	1.501	8.74	54.48	0.255	
	M5	OG	Ap	0–40	0.73	91.03	6.31	2.66	1.531	9.29	1.99	0.215	
			2Ck	40–100	2.28	66.14	31.16	2.70	1.420	9.19	21.94	0.255	
	X		Ap	0–40	9.63 $\pm$ 10.00	87.91 $\pm$ 7.80	6.32 $\pm$ 3.94	5.77 $\pm$ 4.24	1.53 $\pm$ 0.12	9.13 $\pm$ 0.21	6.72 $\pm$ 4.02	0.25 $\pm$ 0.17	
			2Bk/Ck	40–100	12.04 $\pm$ 16.83	64.75 $\pm$ 6.44	15.88 $\pm$ 10.70	19.37 $\pm$ 10.93	1.46 $\pm$ 0.09	8.97 $\pm$ 0.19	28.18 $\pm$ 14.72	0.25 $\pm$ 0.04	
NC-CL <i>n</i> = (3 × 2)	A1	OG	Ap	0–40	1.56	82.93	9.08	7.99	1.570	8.77	9.34	0.271	
			2Ck	40–100	3.80	61.04	16.72	22.24	1.351	8.74	26.16	0.285	
	A2	OG	Ap	0–40	4.28	78.68	8.76	12.56	1.541	8.72	7.80	0.351	
			Ck	40–100	0.64	60.01	22.25	17.74	1.355	8.74	22.76	0.265	
	A3	OG	Ap	0–40	10.85	69.64	12.85	17.51	1.412	8.95	17.13	0.395	
			2Ck	40–100	21.60	51.58	18.83	29.59	1.371	8.83	35.41	0.402	
A4	AT	Ap	0–40	14.47	53.30	25.64	21.06	1.032	8.90	19.41	0.490		
		Ck	40–100	9.83	65.53	14.81	19.66	1.461	8.93	21.26	0.271		
A5	PT	Ap	0–40	3.24	77.06	13.75	9.19	1.370	8.83	9.80	0.560		
		2Bk	40–100	11.99	54.22	18.71	27.07	1.355	8.87	32.37	0.275		
X		Ap	0–40	6.88 $\pm$ 5.51	72.32 $\pm$ 11.67	14.02 $\pm$ 6.87	13.66 $\pm$ 5.54	1.39 $\pm$ 0.22	8.83 $\pm$ 0.09	12.70 $\pm$ 5.21	0.41 $\pm$ 0.11		
		2Bk/2Ck	40–100	9.57 $\pm$ 8.12	58.48 $\pm$ 5.58	18.26 $\pm$ 2.77	23.26 $\pm$ 4.98	1.38 $\pm$ 0.05	8.82 $\pm$ 0.08	27.59 $\pm$ 6.11	0.30 $\pm$ 0.06		

SD \*: Standard deviation; M: Management; SP: Soil sampling; Hor: Horizon; BD: Bulk density; OC: Organic carbon; *n* = Sample size. OC-CL: Old cultivated (tilled > 100 years) Calcisols; NC-CL: Newly cultivated (tilled < 12 years) Calcisols.

In general, the surface horizon (topsoil) was weakly developed with low SOC content ranging from 0.12% (OC-AR: B1-Ap) to 0.56% (NC-CL: PT: A5-Ap). In this sense, the management type (strongly mechanized) caused the soil homogenization by mixing the surface horizons in the first 40 cm. Due to the low SOC, they are not diagnostic horizons [38], so the C content was below the 0.2% (Tables 4 and 5). Similar results were obtained by [54,55] in CL in the Sfax region, near of the study area, stating that the natural SOC content in the Ap horizon of these soils was very low (<0.3%) due to climatic conditions, relief, and lithology. Calcium carbonate content in the deeper horizons (40–100 cm) was variable, with simultaneous presence of secondary precipitation (calcic horizons). According to [56], the soils developed from carbonate parent materials are common and belong to the main soil groups in North Tunisia.

In the remaining soils, the calcium carbonate content was not high, and due to texture (sandy) and the lack of diagnostic horizons, they were classified as AR. The presence of weakly developed AR derived from aeolian sediments has also been confirmed by [37].

Nevertheless, the carbonates' presence in these soils was expressed using the Calcaric qualifier (C1, C2, B1, B2).

### 3.2. Soil Bulk Density and pH in AR and CL

One of the problems that the soils in the study area have is bulk density (BD) quantification, since they are sandy soils with strong wind erosion, together with the high carbonate concentration. In this sense, the BD determination is very important, since it affects SOCS analysis [50–52].

The BD study did not show a clear trend; although in most cases BD decreased in depth, this behavior was expected, since in most cases, the BD reduction prevented an increase in the SOC concentration (Tables 4 and 5). In the superficial layer (0–40 cm), our results showed that BD on average were between 1.7 and 1.39 Mg m<sup>-3</sup> for OC-AR and NC-CL, respectively; however, in sub-soil (40–100), the BD ranged between 1.69 and 1.35 Mg m<sup>-3</sup> for NC-AR and NC-CL. It is important to note that BD was higher in AR compared to CL regardless of the study period (long-term and short-term) (Tables 4 and 5). The studied soils have shown that BD decreased significantly in NC-CL and OC-CL; in this line, [57] confirmed that in temperate zones, a decrease in BD is related to an increase in calcium carbonate content. BD values tended to be higher in surface horizons than in deep layers specifically in CL. This increase can be explained by the effect of tillage, which reduces the SOM content, increasing soil compaction and increasing the soil DB. This effect has been shown by [58] in soils with different land uses, indicating a BD increase in cropland. In addition, this BD increase was more important in the soil that was most intensively tilled.

Our results show differences in BD between CL and AR, and these differences may be due to soil type, management practices, and climatic conditions, in addition due to several factors, such as volume and rainfall intensity, drying and wetting of soil, land position, and crop type, among others [59]. According to [60], the time factor could be the most important factor that explains the BD variations in surface and subsurface. In fact, these authors reported that under conventional tillage, the general trend in surface at 15 cm depth was a BD increase during the cropping cycle except in the surface rows where BD was already high at the first measurement time.

The soil pH values showed that pH decreased slightly with increasing depth in newly and previously cultivated CL and AR (Tables 4 and 5). In the superficial layer, pH values were between 8.78 and 8.95 in NC-CL and between 8.94 and 9.17 in NC-AR. In the deep layer, pH values ranged from 8.74 to 8.93 in NC-CL and from 8.81 to 9.12 in NC-AR. In OC soil, pH values in superficial layers fluctuated between 8.78 and 9.29 in OC-CL and between 9.10 to 9.25 in OC-AR. In the deep layer, pH values varied between 8.74 and 9.19 in OC-CL and between 9.10 and 9.19 in OC-AR. These results were similar to those found by [61] in Sfax Tunisia in conventional tillage (cultivated soil) with alkaline pH ( $9.17 \pm 0.76$ ); since the SOC degradation and the CaCO<sub>3</sub> solubilization may affect pH, in this line, we suggest that the alkaline condition can be caused by the high concentration of the calcium carbonate.

### 3.3. Soil Organic Carbon Stock Concentration

All SOCS values of newly cultivated soils (<12 years) and old cultivated soils (>100 years) are shown in Tables 6 and 7. The results indicate important differences between SOCS in newly and previously cultivated soils regarding the soil type (CL or AR).



Table 6. Soil organic carbon stock in Arenosols.

Soil	M	SP	LU	Hor	Th (cm)	Gravel (%)	BD (Mg m <sup>-3</sup> )	OC (g kg <sup>-1</sup> )	SOCS Mg ha <sup>-1</sup>	T-SOCS Mg ha <sup>-1</sup>
AR	OC-AR <i>n</i> = (3 × 2)	B1	OG	Ap	40	1.15	1.701	1.25	8.407	40.783
				AC	60	0.60	1.645	3.30	32.376	
		B2	OG	Ap	40	1.91	1.692	2.90	19.252	35.260
				C	60	6.74	1.673	1.71	16.008	
		B3	OG	Ap	40	1.48	1.705	2.82	18.947	47.436
				AC	60	2.60	1.681	2.90	28.489	
	X		Ap	40	1.51 ± 0.28	1.70 ± 0.01	2.32 ± 0.93	15.535 ± 6.175	41.159 ± 7.363	
			AC	60	3.31 ± 3.13	1.67 ± 0.02	2.64 ± 0.83	25.624 ± 8.552		
	NC-AR <i>n</i> = (3 × 2)	C1	AT	Ap	40	2.98	1.560	2.70	16.346	30.839
				C	60	4.32	1.530	1.65	14.493	
		C2	OG	Ap	40	0.66	1.632	2.90	18.806	28.303
				BC	60	0.88	1.681	0.95	9.497	
		C3	OG	Ap	40	0.09	1.670	1.45	9.677	16.239
				C	60	0.44	1.690	0.65	6.562	
X		Ap	40	1.24 ± 1.53	1.62 ± 0.06	2.35 ± 0.79	14.943 ± 4.723	25.127 ± 4.367		
		C	60	1.88 ± 2.13	1.63 ± 0.09	1.08 ± 0.51	10.184 ± 4.010			

M: Management; SP: Soil sampling; BD: Bulk density; OC: Organic carbon; SOCS: Soil organic carbon stock; T-SOCS: Total SOCS; *n* = Sample size. OC-CL: Old cultivated (tilled > 100 years) Calcisols; NC-CL: Newly cultivated (tilled < 12 years) Calcisols.

Table 7. Soil organic carbon stock in Calcisol.

Soil	M	SP	LU	Hor	Th (cm)	Gravel (%)	BD (Mg m <sup>-3</sup> )	OC (g kg <sup>-1</sup> )	SOCS Mg ha <sup>-1</sup>	T-SOCS Mg ha <sup>-1</sup>
CL	OC-CL <i>n</i> = (3 × 2)	M1	OG	Ap	40	6.91	1.621	1.31	7.907	25.661
				2Ck	60	7.47	1.591	2.01	17.754	
		M2	OG	Ap	40	0.81	1.610	2.60	16.608	39.796
				2Bk	60	2.38	1.342	2.95	23.188	
		M3	OG	Ap	40	23.54	1.330	5.40	21.965	33.607
				Ck	60	41.86	1.451	2.30	11.642	
	M4	OG	Ap	40	16.16	1.570	1.15	6.055	27.596	
			2Ck	60	6.20	1.501	2.55	21.541		
	M5	OG	Ap	40	0.73	1.531	2.15	13.070	34.301	
			2Ck	60	2.28	1.420	2.55	21.231		
	X		Ap	40	9.63 ± 10.00	1.53 ± 0.12	2.52 ± 1.72	13.121 ± 6.471	32.192 ± 5.536	
			2Bk/Ck	60	12.04 ± 16.83	1.46 ± 0.09	2.47 ± 0.35	19.071 ± 4.600		
	NC-CL <i>n</i> = (3 × 2)	A1	OG	Ap	40	1.56	1.570	2.71	16.753	38.977
				2Ck	60	3.80	1.351	2.85	22.224	
A2		OG	Ap	40	4.28	1.541	3.51	20.710	42.117	
			Ck	60	0.64	1.355	2.65	21.407		
A3		OG	Ap	40	10.85	1.412	3.95	19.889	45.815	
			2Ck	60	21.60	1.371	4.02	25.926		
A4	AT	Ap	40	14.47	1.032	4.90	17.300	38.721		
		Ck	60	9.83	1.461	2.71	21.421			
A5	PT	Ap	40	3.24	1.370	5.60	29.694	49.371		
		2Bk	60	11.99	1.355	2.75	19.677			
X		Ap	40	6.88 ± 5.51	1.39 ± 0.22	4.13 ± 1.14	20.869 ± 5.209	43.000 ± 3.7625		
		2Bk/2Ck	60	9.57 ± 8.12	1.38 ± 0.05	3.00 ± 0.58	22.131 ± 2.316			

M: Management; SP: Soil sampling; BD: Bulk density; OC: Organic carbon; SOCS: Soil organic carbon stock; T-SOCS: Total SOCS; *n* = Sample size. OC-CL: Old cultivated (tilled > 100 years) Calcisols; NC-CL: Newly cultivated (tilled < 12 years) Calcisols.

The SOCS analysis showed that on average, the SOCS content in CL was 20.9 Mg ha<sup>-1</sup> and 13.1 Mg ha<sup>-1</sup> in topsoil (0–40 cm depth) for NC-CL and OC-CL, respectively. In sub-soil (1 m depth), the SOCS was 43.0 Mg ha<sup>-1</sup> (NC-CL) and 32.2 Mg ha<sup>-1</sup> (OC-CL). In this line, [62] in dryland of southeastern Spain found 52 Mg ha<sup>-1</sup> and 70 Mg ha<sup>-1</sup> for 0.5 m

and 1 m depth, respectively, and [63] in southern Spain reported 50.1 Mg ha<sup>-1</sup> at 0.25 m depth and 82 Mg ha<sup>-1</sup> at 0.75 m depth. These differences in SOCS between our results and the results reported by [62,63] could be explained by the impact of tillage that increases the SOC depletion and reduces the biomass generation. In addition, these discrepancies (low SOCS content in the studied soils) may reflect a combination of different factors such as intensive farming practices, climate conditions, soil types, soil age, and topography area [64,65].

In the AR case, on average, the SOCS contents were as follows: 14.9 Mg ha<sup>-1</sup> and 25.1 Mg ha<sup>-1</sup> for 0.4 m and 1 m depth, respectively, in NC-AR, and 15.5 Mg ha<sup>-1</sup> and 41.2 Mg ha<sup>-1</sup> at 0.4 m and 1 m depth, respectively, in OC-AR. Other authors such as [66], for soil groups in Peninsular Spain (FAO soil map of Peninsular Spain), reported low SOCS content in AR (22.2 Mg ha<sup>-1</sup>) in soil profiles deeper than 1 m. Furthermore, [8] indicated that the SOC content (average values) in the upper 1 m was 31 Mg ha<sup>-1</sup> for sandy AR. These differences between our data and the data indicated by other authors may be due to soil texture (sandy soils) and tillage. In NC-AR, the SOCS at surface depth (Ap horizon) was higher than in the C horizon. In this line [67], argued that the SOCS on surface horizon was greater than in deep soil due to tillage and in turn increasing the physical protection of native SOC from microbial decomposition. We can increase C inputs into surface soil by enhancing crop biomass and in turn residue return. With respect to OC-AR, this relation was inverse (SOCS increased in AC horizon); this effect can be explained by soil texture (sandy soils) and tillage, because native SOC can be reduced on the surface, which may be attributed to soluble organic compounds that can leach into deeper layers, increasing the soil aggregates [68,69].

### 3.4. Effects of Management Time on Soil Organic Carbon Stock

The results indicated important differences between SOCS (0–100 cm) in newly and previously cultivated soils regarding the soil type (CL or AR). In NC-CL, the SOCS (0–100 cm) content ranged from 38.7 Mg ha<sup>-1</sup> (NL-CL-AT) to 49.4 Mg ha<sup>-1</sup>, (NC-CL-PT); however, lower SOCS were obtained for OC-CL, varying between 25.7 Mg ha<sup>-1</sup> (OC-CL-OG) and 39.8 Mg ha<sup>-1</sup> (OC-CL-OG). In the CL case, it is important to note that the land use affected the SOCS content, since on average (0–100 cm), the centenary OG (OC-CL) had a 25.6% lower SOCS than the young OG (NC-CL). However, in depth (40–100 cm), no differences were found according to the land use age (OC and NC) with respect to SOCS. Another aspect to highlight is that in the different crops developed (olive, almond, and pistachio) in NC-CL, the pistachio (PT) had the highest SOCS content (49.4 Mg ha<sup>-1</sup>), whereas the lowest SOCS values were found in almond tree (AT) (38.7 Mg ha<sup>-1</sup>).

However, the study of AR has shown that the SOCS content on average in the Ap horizon (0–40 cm) in NC and OC were very similar (OC-AR: 15.5 Mg ha<sup>-1</sup>; NC-AR: 14.9 Mg ha<sup>-1</sup>), but nevertheless, in depth (40–100 cm: AC/BC/C horizon), significant differences were found (OC-AR: 25.6 Mg ha<sup>-1</sup>; NR-AR: 10.2 Mg ha<sup>-1</sup>). When comparing both soils (AR and CL), significant differences were observed between NC-CL and NC-AR; however, no significant differences were observed between OC-CL and OC-AR. The highest SOCS contents were obtained in NC-CL regardless of the land tillage time.

In OC-CL, the SOCS (32.2 Mg ha<sup>-1</sup>) was significantly lower compared to NC-CL (43 Mg ha<sup>-1</sup>). However, the SOCS in OC-AR (41.2 Mg ha<sup>-1</sup>) increased with respect to NC-AR (25.1 Mg ha<sup>-1</sup>) (Tables 6 and 7). These results indicate that the SOCS in CL and AR are linked to land management duration (OC > 100 years and NC < 12 years). In this line, our results are accordance with [70] in CL in Turkey, which justified an SOC reduction with intensive soil management under strong wind and water erosion conditions, clarifying that under these conditions, the SOM is rapidly mineralized in the soil. In addition, soil management over time may deteriorate the aggregates' stability [69] and reduce the CL quality, affecting the soil water storage capacity in rainfed OG [70], these processes therefore could explain the SOM reduction and the SOCS decreasing in the surface horizon.

Several studies have found an increase in the SOC content in AR cultivated more than 20 years in different arid areas [71–73]. This is in concordance with our study; however, according to [74], the SOC could be depleted under intensive tillage, which is in discrepancy with our findings. In this sense, [75], in the same study area as [76], showed the olive roots' presence within the sands; however, it was observed that olive roots were absent in the calcareous crusts. In the study soils, more than 90% of AR had sandy texture; for this reason, we suggest that the SOC increase could be due to the olive roots' presence since the calcareous crusts are not present. Moreover, the tillage can affect a large part in the accumulation of the olive roots (along the profile), which may explain our findings. In the literature, the tillage is assumed to alter the SOCS and also to reduce the SOC quantity and to deteriorate the aggregates in the soil [10,69]. However, long cultivation periods with heavy tillage showed a positive effect on SOCS in AR, but due to the climatic conditions (semi-desert), more research should be carried out in AR to be able to identify the mechanism that explains this SOCS increase in AR.

Another question to highlight is that OC could be influenced by the abundance of the olive root mass. In this sense, it should be noted that the olive tree renews its roots every year and that the root system is better developed in the ventilated area (AR) than in compacted soils (CL). Another issue to point out is the addition of the organic amendments, so that some sources of organic amendments may accelerate the SOC mineralization. In the study area, in some plots, different organic amendments were added (B1, A1, A2, and M1), the main consequence was a SOCS reduction in the soil surface with respect to the plots without amendments (C1, A4, A5, and M3) (Tables 2 and 3). However, in other cases, there was a SOCS increase in the soil surface in modified plots (A3, B2, C2, and M2) compared to the unmodified plots (M4). This SOCS variability could be explained by the results obtained by [76], who suggested that depending on the organic amendment type, the SOC balance (gain or loss of SOC) is conditioned by the incubation period and the organic amendment type. The increase in SOC of green manure derived from olive pruning residues (dried and crushed shoots and leaves of olive trees) is too limited ( $0.13 \text{ mg g}^{-1}$ ), and the SOC balance (between the day after amendment and 120 days after) ( $-0.45$ ) compared to the SOC gain of compost of manure and olive husk and palm-leaf-based compost are ( $0.28 \text{ mg g}^{-1}$ ) and ( $0.31 \text{ mg g}^{-1}$ ), respectively. The SOC balance is ( $-0.02$ ) and ( $-0.05$ ) in both compost types. These results could explain the SOC variability in the study area. However, we must be careful, as the organic amendments addition can be confounding as they lead to variability of the SOC content.

In Egypt (semi-desert conditions), Ref. [77] proposed a direct relation between the cropping history and SOCS (longer cultivation period implies the higher SOCS). This result is in agreement with AR and in disagreement with CL. We suggest that the soil type plays an important role in the depletion or the gain of SOC. In the study area, AR are efficient for SOCS accumulation, especially after 100 years of cultivation and tillage, and they could be carbon sink and may be involved in reducing carbon dioxide ( $\text{CO}_2$ ) emissions. In fact, it has been reported that soil might represent a sink for atmospheric  $\text{CO}_2$  [78–80].

### 3.5. Uncertainties and Bias

The complexity of these soils study is due to the continuous movement of the surface particles by the wind (eolian processes). For this reason, in demi-desert areas, few studies have been developed to define SOCS thresholds. In this line, all soil samples were taken at the same time (synchronic approach) under different management practices at known durations from an initial reference state, and the SOCS was compared under this initial reference state [53]. However, due to eolian processes (displacement of particles by the wind), it could be useful to carry out different samplings throughout the year, to reduce the wind effect, this question is relevant to establish the thickness of the surface horizon.

However, despite the mistakes, the SOCS study in these soils type (AR and CL) under these climatic conditions could help us to promote strategies to combat the desertifica-

tion and climate change. So, this research constitutes a preliminary assessment of SOCS estimation with different land uses and different management practice durations.

#### 4. Conclusions

The main conclusions derived from this research depend on land use (olive grove, almond tree, and pistachio tree), land management (tillage and amendments) the practice's duration (old plantations > 100 years and new plantations < 12 years) and soil type (Arenosols and Calcisols) in semi-desert conditions. In this line, the crop type, tillage, and tillage duration affect to soil bulk density, pH, and SOC for the same soil type.

On average, the SOCS for woody crops in semi-desert areas in CL is  $43.0 \text{ Mg ha}^{-1}$  and  $32.2 \text{ Mg ha}^{-1}$  (0–100 cm depth) for NC and OC, respectively, and with 40 cm depth, the SOCS is  $20.9 \text{ Mg ha}^{-1}$  (NC) and  $13.2 \text{ Mg ha}^{-1}$  (OC). In the case of the AR, the SOCS is  $25.1 \text{ Mg ha}^{-1}$  and  $41.2 \text{ Mg ha}^{-1}$  (0–100 cm depth) for NC and OC, respectively, and with 40 cm depth, the SOCS is  $14.9 \text{ Mg ha}^{-1}$  (NC) and  $15.5 \text{ Mg ha}^{-1}$  (OC). According to this, in dryland (semi-desert conditions), some soils could have a good capacity to increase soil organic carbon with certain management practices and duration specifically, the AR could increase the SOCS after 100 years of cultivation and tillage; however, the CL the SOC can be reduced.

Thus, SOCS content variations were detected and have established pistachio as the most efficient woody crop related to carbon storage ( $49.4 \text{ Mg ha}^{-1}$ ) under the soils and climatic conditions analyzed in the short term. Therefore, good information to better understand the dynamics of soil organic carbon storage could help to develop and consolidate the Framework of the United Nation Convention to Combat Desertification (UNCCD) on the one hand and to minimize greenhouse gases on the other.

**Author Contributions:** Conceptualization, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., methodology, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., software, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., validation, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., formal analysis, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., investigation, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., data curation, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., writing—original draft preparation, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., writing—review and editing, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., visualization, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B., supervision, F.B., R.C., H.R., H.B.M., K.G., N.R., N.B., M.S., M.G.-R., L.P.-A., and Á.S.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** The Ministry of Agriculture and Water Resources and the Ministry of Higher Education and Scientific Research (Tunisia) supported this study.

**Institutional Review Board Statement:** Not applicable.

**Data Availability Statement:** The data that support this study cannot be publicly shared due to ethical or privacy reasons and may be shared upon reasonable request to the corresponding author if appropriate.

**Acknowledgments:** The study design and data collection were carried out at the Research Laboratory of Sustainable Olive and Arboriculture Growing of Olive Institute (Tunisia). Analysis of soil samples were carried out at the Department of Evolutionary Biology, Ecology and Environmental Science of the University of Barcelona. The facilities and services of the Olive Institute of Sfax (Tunisia), NCU in Toruń (Poland), and the Universities of Barcelona, Cadiz, and Cordoba (Spain) are gratefully acknowledged. The authors thank Nabil Soua (Olive) Institute for his expertise, hard work and all contribute to the field and in the laboratory. The authors are grateful to the members, specifically Mouhamed Fakhfekh, of the organic farm located in the Menzel Chaker region in Sfax, Tunisia. We are grateful to Souha Hammouda for her support and for her assistance in improving the English quality of the manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Lal, R. Carbon sequestration. *Philos. Trans. R. Soc. Lond.* **2008**, *63*, 815–830. [[CrossRef](#)] [[PubMed](#)]
2. Stockmann, U.; Adams, M.A.; Crawford, J.W.; Field, D.J.; Henakaarchchi, N.; Jenkins, M.; Minasny, B.; McBratney, A.B.; Courcelles, V.; Singh, K.; et al. The knowns, known unknowns and unknowns of sequestration of soil organic carbon. *Agric. Ecosyst. Environ.* **2013**, *164*, 80–99. [[CrossRef](#)]
3. Brahim, N.; Ibrahim, H. Effect of Land Use on Organic Carbon Distribution in a North African Region: Tunisia Case Study. In *Soil Management and Climate Change*; Academic Press: Cambridge, MA, USA, 2018; pp. 15–24. [[CrossRef](#)]
4. Pansu, M.; Ibrahim, H.; Hatira, A.; Brahim, N.; Drevon, J.J.; Harmand, J.M.; Chotte, J.L.; Blavet, D. Modelling the continuous exchange of nitrogen between microbial decomposers, the organs and symbionts of plants, soil reserves and the atmosphere. *Soil Biol. Biochem.* **2018**, *125*, 185–196. [[CrossRef](#)]
5. Lal, R.; Smith, P.; Jungkunst, H.F.; Mitsch, W.J.; Lehmann, J.; Nair, P.K.R.; McBratney, A.B.; de Moraes Sá, J.C.; Schneider, J.; Zinn, Y.L.; et al. The carbon sequestration potential of terrestrial ecosystems. *J. Soil Water Conserv.* **2018**, *73*, 145A–152A. [[CrossRef](#)]
6. Lefevre, C.; Rekik, F.; Alcantara, V.; Wiese, L. *Soil Organic Carbon: The Hidden Potential*; FAO: Rome, Italy, 2017.
7. Salcedo, I.H.; Sampaio, V.S.B. Matéria orgânica do solo no Bioma Caatinga. In *Fundamentos da Matéria orgânica do Solo, Ecossistemas tropicais e Subtropicais*; Santos, G., de Silva, A.L.S., da Canellas, L.P., de Camargo, F.A.O., Eds.; Metrópole: Porto Alegre, Brasil, 2008; Volume 22, pp. 419–441.
8. Batjes, N.H. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* **1996**, *47*, 151–163. [[CrossRef](#)]
9. Mörchen, R.; Lehdorff, E.; Diaz, F.A.; Moradi, G.; Bol, R.; Fuentes, B.; Klumpp, E.; Amelung, W. Carbon accrual in the Atacama Desert. *Glob. Planet. Chang.* **2019**, *181*, 102993. [[CrossRef](#)]
10. González-Rosado, M.; Parras-Alcántara, L.; Aguilera-Huertas, J.; Lozano-García, B. Long-term evaluation of the initiative 4% under different soil managements in Mediterranean olive groves. *Sci. Total Environ.* **2021**, *758*, 143591. [[CrossRef](#)] [[PubMed](#)]
11. Conant, R.T.; Ryan, M.G.; Ågren, G.I.; Birge, H.E.; Davidson, E.A.; Eliasson, P.E.; Evans, S.E.; Frey, S.D.; Giardina, C.P.; Hopkins, F.M.; et al. Temperature and soil organic matter decomposition rates—Synthesis of current knowledge and a way forward. *Glob. Chang. Biol.* **2011**, *17*, 3392–3404. [[CrossRef](#)]
12. Hammad, H.M.; Khaliq, A.; Abbas, F.; Farhad, W.; Fahad, S.; Aslam, M.; Bakhat, H.F. Comparative effects of organic and inorganic fertilizers on soil organic carbon and wheat productivity under arid region. *Commun. Soil Sci. Plant Anal.* **2020**, *51*, 1406–1422. [[CrossRef](#)]
13. Gargouri, K.; Rigane, H.; Arous, I.; Touil, F. Evolution of soil organic carbon in an olive orchard under arid climate. *Sci. Hortic.* **2013**, *152*, 102–108. [[CrossRef](#)]
14. González-Rosado, M.; Lozano-García, B.; Aguilera-Huertas, J.; Parras-Alcántara, L. Short-term effects of land management change linked to cover crop on soil organic carbon in Mediterranean olive grove hillsides. *Sci. Total Environ.* **2020**, *744*, 140683. [[CrossRef](#)] [[PubMed](#)]
15. Noy-Meir, I. Desert Ecosystems: Environment and Producers. *Ann. Rev. Ecol. Syst.* **1973**, *4*, 25–51. [[CrossRef](#)]
16. Campbell, A.; Miles, L.; Lysenko, I.; Huges, A.; Gibbs, H. *Carbon Storage in Protected Areas*; Technical Report; UNEP World Conservation Monitoring Center: Cambridge, UK, 2008.
17. Pointing, S.B.; Belnap, J. Microbial Colonization and Controls in Dryland Systems. *Nat. Rev. Microbiol.* **2012**, *10*, 551–562. [[CrossRef](#)] [[PubMed](#)]
18. Koyama, A.; Harlow, B.; Evans, R.D. Greater Soil Carbon and Nitrogen in a Mojave Desert Ecosystem after 10 Years Exposure to Elevated CO<sub>2</sub>. *Geoderma* **2019**, *355*, 113915. [[CrossRef](#)]
19. Zhao, K.; Kong, W.; Wang, F.; Long, X.E.; Guo, C.; Yue, L. Desert and Steppe Soils Exhibit Lower Autotrophic Microbial Abundance but Higher Atmospheric CO<sub>2</sub> Fixation Capacity Than Meadow Soils. *Soil Biol. Biochem.* **2018**, *127*, 230–238. [[CrossRef](#)]
20. Brahim, N.; Ibrahim, H.; Hatira, A. Tunisian Soil Organic Carbon Stock—Spatial and Vertical Variation. *Procedia Eng.* **2014**, *69*, 1549–1555. [[CrossRef](#)]
21. Bayer, C.; Martin-Neto, L.; Mielniczuk, J.; Pavinato, A.; Diekow, J. Carbon sequestration in two Brazilian Cerrado soils under no-till. *Soil Tillage Res.* **2006**, *86*, 237–245. [[CrossRef](#)]
22. FAO. *Organizing Committee of the First Symposium on the Management of Tropical Sandy Soils through the Land Management Officer*; FAO Regional Office for Asia and the Pacific: Bangkok, Thailand, 2005.
23. Azlan, A.; Aweng, E.R.; Ibrahim, C.O.; Noorhaidah, A. Correlation between Soil Organic Matter, Total Organic Matter and Water Content with Climate and Depths of Soil at Different Land use in Kelantan. *Malays. J. Appl. Sci. Environ. Manag.* **2012**, *16*, 353–358.
24. Ding, Z.; Kheir, A.M.; Ali, M.G.; Ali, O.A.; Abdelaal, A.I.; Zhou, Z.; He, Z. The integrated effect of salinity, organic amendments, phosphorus fertilizers, and deficit irrigation on soil properties, phosphorus fractionation and wheat productivity. *Sci. Rep.* **2020**, *10*, 1–13. [[CrossRef](#)]
25. Ben Mbarek, H.; Gargouri, K.; Mbadra, C.; Ben Mahmoud, I.; Chaker, R.; Maktouf, S.; Abbas, O.; Baeten, V.; Rigane, H. Effects of combination of tillage with olive mill wastewater on soil organic carbon groups in arid soils. *Arab. J. Geosci.* **2020**, *13*, 255. [[CrossRef](#)]
26. Park, C.E.; Jeong, S.J.; Joshi, M.; Osborn, T.J.; Ho, C.H.; Piao, S.; Feng, S. Keeping global warming within 1.5 C constrains emergence of aridification. *Nat. Clim. Chang.* **2018**, *8*, 70–74. [[CrossRef](#)]
27. Ma, X.; Zhu, J.; Yan, W.; Zhao, C. Projections of desertification trends in Central Asia under global warming scenarios. *Sci. Total Environ.* **2021**, *781*, 146777. [[CrossRef](#)] [[PubMed](#)]

28. Moussadek, R.; Mrabet, R.; Dahan, R.; Zouahri, A.; El Mourid, M.; Ranst, E.V. Tillage System Affects Soil Organic Carbon Storage and Quality in Central Morocco. *Appl. Environ. Soil Sci.* **2014**, 1–8. [[CrossRef](#)]
29. Jaouadi, J.; Brahim, N.; Hatira, A.; Aichi, H.; Ben Hassine, H. Distribution of organic carbon stocks in cambisol under two different crops in semi-arid Tunisian climate. *J. Res. Environ. Earth Sci.* **2019**, *8*, 216–222.
30. Znaïdi, A.; Brahim, N.; Ibrahim, H.; Bol, R.; Chaouachi, M. Comparison of organic carbon stock of Regosols under two different climates and land use in Tunisia. *Arab. J. Geosci.* **2020**, *13*, 1011. [[CrossRef](#)]
31. Macías, M.D.J.G.; Carbajal, N.; Vargas, J.T. Soil deterioration in the southern Chihuahuan Desert caused by agricultural practices and meteorological events. *J. Arid Environ.* **2020**, *176*, 104097. [[CrossRef](#)]
32. Fallahzade, J.; Karimi, A.; Naderi, M.; Shirani, H. Soil mechanical properties and wind erosion following conversion of desert to irrigated croplands in central Iran. *Soil Tillage Res.* **2020**, *204*, 104665. [[CrossRef](#)]
33. González-Rosado, M.; Parras-Alcántara, L.; Aguilera-Huertas, J.; Lozano-García, B. Soil Productivity Degradation in a Long-Term Eroded Olive Orchard under Semiarid Mediterranean Conditions. *Agronomy* **2021**, *11*, 812. [[CrossRef](#)]
34. Hu, Y.; Wang, Z.; Zhang, Z.; Song, N.; Zhou, H.; Li, Y.; Hale, L. Alteration of desert soil microbial community structure in response to agricultural reclamation and abandonment. *Catena* **2021**, *207*, 105678. [[CrossRef](#)]
35. Köberl, M.; Müller, H.; Ramadan, E.M.; Berg, G. Desert farming benefits from microbial potential in arid soils and promotes diversity and plant health. *PLoS ONE* **2011**, *6*, e24452. [[CrossRef](#)]
36. Kottek, M.; Grieser, J.; Beck, C.; Rudolf, B.; Rubel, F. World map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* **2006**, *15*, 259–263. [[CrossRef](#)]
37. Previtali, F.; Gasmı, N.; Julitta, F.; Cantelli, D.; Deroma, M.; Madrau, S.; Zucca, C. The geomorphologic and pedo-morphologic maps of Skhirat (Feriana, Tunisia) at the 1:50000 scale. *J. Maps* **2014**, *10*, 73–81. [[CrossRef](#)]
38. IUSS Working Group WRB. World Reference Base for soil resources 2014, update 2015. In *International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*; World Soil Resources Report No. 106; FAO: Rome, Italy, 2015.
39. Driessen, P.M.; Deckers, J.; Spaargaren, O. *Lecture Notes on the Major Soils of the World*; World Soil Resources Reports; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 2001; Volume 94.
40. Missaoui, A.; Zerai, K.; Feridhi, A.; Baroudi, M.; Baroudi, J. *Atlas du Gouvernourat de Sfax*; Direction Générale de l'aménagement du territoire: Tunisia, Tunisia, 2013; p. 105.
41. Lal, R. Forest soils and carbon sequestration. *For. Ecol. Manag.* **2005**, *220*, 242–258. [[CrossRef](#)]
42. Parras-Alcántara, L.; Lozano-García, B.; Brevik, E.C.; Cerdà, A. Soil organic carbon stocks assessment in Mediterranean natural areas: A comparison of entire soil profiles and soil control sections. *J. Environ. Manag.* **2015**, *155*, 219–228. [[CrossRef](#)] [[PubMed](#)]
43. Álvaro-Fuentes, J.; Lóczy, D.; Thiele-Bruhn, S.; Zornoza, R. *Handbook of Plant and Soil Analysis for Agricultural Systems*; Crai uptc Editions: Cartagena, Spain, 2019; p. 389.
44. Blake, G.R.; Hartge, K.H. Bulk density. In *Methods of Soil Analysis, Part 1: Physical and Mineralogical Methods*, 2nd ed.; Klute, Ed.; ASA, SSSA: Madison, WI, USA, 1986; pp. 377–382.
45. USDA. *Soil Survey Laboratory Methods Manual, Soil Survey Investigation Report no. 42, Version 4.0*; USDA-NCRS: Lincoln, NE, USA, 2004.
46. Guitián, F.; Carballas, T. *Técnicas de Análisis de Suelos*; Picro Sacro: Santiago de Compostela, Spain, 1976.
47. Nelson, D.W.; Sommers, L.E. Total carbon, organic carbon and organic matter. In *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties. Agronomy Monograph*; Page, A.L., Miller, R.H., Keeney, D., Eds.; ASA and SSSA: Madison, WI, USA, 1982; Volume 9, pp. 539–579.
48. FAO. Standard operating procedure for soil calcium carbonate equivalent. In *Volumetric Calcimeter Method*; FAO: Roma, Italy, 2020.
49. Estefan, G.; Sommer, R.; Ryan, J. *Methods of Soil, Plant, and Water Analysis-A Manual for the West Asia and North Africa Region*; MEL: Vermont, VI, Australia, 2013; pp. 170–176.
50. Aguilera-Huertas, J.; Lozano-García, B.; González-Rosado, M.; Parras-Alcántara, L. Effects of Management and Hillside Position on Soil Organic Carbon Stratification in Mediterranean Centenary Olive Grove. *Agronomy* **2021**, *11*, 650. [[CrossRef](#)]
51. Stolbovoy, V.; Montanarella, L.; Filippi, N.; Jones, A.; Gallego, J.; Grassi, G. *Soil Sampling Protocol to Certify the Changes of Organic Carbon Stock in Mineral Soil of the European Union*; Version 2; EUR 21576 EN/2. 56; Office for Official Publications of the European Communities: Luxembourg, 2007; ISBN 978-92-79-05379.
52. IPCC. Intergovernmental Panel on Climate Change. In *Good Practice Guidance for Land Use, Land Use Change and Forestry*; Penman, J., Gytarsky, M., Hiraishi, T., Krug, T., Kruger, D., Pipatti, R., Buendia, L., Miwa, K., Ngara, T., Tanabe, K., Eds.; IPCC/OECD/IEA/IGES: Hayama, Japan, 2003.
53. Costa Junior, C.; Corbeels, M.; Bernoux, M.; Piccolo, M.C.; Siqueira Neto, M.; Feigl, B.J.; Cerri, C.E.P.; Cerri, C.C.; Scopel, E.; Lal, R. Assessing soil carbon storage rates under no-tillage: Comparing the synchronic and diachronic approaches. *Soil Tillage Res.* **2013**, *134*, 207–212. [[CrossRef](#)]
54. Belaid, N.; Feki, S.; Cheknane, B.; Frank-Neel, C.; Baudu, M. Impacts of irrigation systems on vertical and lateral metals distribution in soils irrigated with treated wastewater: Case study of Elhajeb-Sfax. *Agric. Water Manag.* **2019**, *225*, 105739. [[CrossRef](#)]
55. Belaid, N.; Neel, C.; Kallel, M.; Ayoub, T.; Ayadi, A.; Baudu, M. Long term effects of treated wastewater irrigation on calcisol fertility: A case study of Sfax-Tunisia. *Agric. Sci.* **2012**, *3*, 702–713. [[CrossRef](#)]

56. Dewitte, O.; Jones, A.; Spaargaren, O.; Breuning-Madsen, H.; Brossard, M.; Dampha, A.; Deckers, J.; Gallali, T.; Hallett, S.; Jones, R.; et al. Harmonisation of the soil map of Africa at the continental scale. *Geoderma* **2013**, *211*, 138–153. [[CrossRef](#)]
57. Świtoniak, M.; Dąbrowski, M.; Łyszkiewicz, A. The Influence of Human-induced Erosion on the Soil Organic Carbon Stock in Vineyards of Fordon Valley. *Pol. J. Soil Sci.* **2015**, *48*, 197. [[CrossRef](#)]
58. Bormann, H.; Klaassen, K. Seasonal and land use dependent variability of soil hydraulic and soil hydrological properties of two Northern German soils. *Geoderma* **2008**, *145*, 295–302. [[CrossRef](#)]
59. Ordoñez-Morales, K.; Cadena-Zapata, M.; Zermeño-González, A.; Campos-Magaña, S. Effect of Tillage Systems on Physical Properties of a Clay Loam Soil under Oats. *Agriculture* **2019**, *9*, 62. [[CrossRef](#)]
60. Alletto, L.; Coquet, Y. Temporal and spatial variability of soil bulk density and near-saturated hydraulic conductivity under two contrasted tillage management systems. *Geoderma* **2009**, *152*, 85–94. [[CrossRef](#)]
61. Ben Mbarek, H.; Gargouri, K.; Mbadra, C.; Chaker, R.; Souidi, Y.; Abbas, O.; Baeten, V.; Rigane, H. Change and spatial variability of soil organic matter humification after long-term tillage and olive mill wastewater application in arid regions. *Soil Res.* **2020**, *58*, 388. [[CrossRef](#)]
62. Díaz-Hernández, J.L.; Fernández, E.B.; González, J.L. Organic and inorganic carbon in soils of semiarid regions: A case study from the Guadix–Baza basin (Southeast Spain). *Geoderma* **2003**, *114*, 65–80. [[CrossRef](#)]
63. Muñoz-Rojas, M.; De la Rosa, D.; Zavala, L.M.; Jordán, A.; Anaya-Romero, M. Changes in land cover and vegetation carbon stocks in Andalusia, Southern Spain (1956–2007). *Sci. Total Environ.* **2011**, *409*, 2796–2806. [[CrossRef](#)] [[PubMed](#)]
64. Hu, T.; Sørensen, P.; Olesen, J.E. Soil carbon varies between different organic and conventional management schemes in arable agriculture. *Eur. J. Agron.* **2018**, *94*, 79–88. [[CrossRef](#)]
65. Yigini, Y.; Panagos, P. Assessment of soil organic carbon stocks under future climate and land cover changes in Europe. *Sci. Total Environ.* **2016**, *557*, 838–850. [[CrossRef](#)] [[PubMed](#)]
66. Rodríguez-Murillo, J.C. Organic carbon content under different types of land use and soil in peninsular Spain. *Biol. Fertil. Soils* **2001**, *33*, 53–61. [[CrossRef](#)]
67. Blanco-Canqui, H.; Lal, R. No-tillage and soil-profile carbon sequestration: An on farm assessment. *Soil Sci. Soc. Am. J.* **2008**, *72*, 693–701. [[CrossRef](#)]
68. Diekow, J.; Mielniczuk, J.; Knicker, H.; Bayer, C.; Dick, D.P.; Kögel-Knabner, I. Soil C and N stocks as affected by cropping systems and nitrogen fertilisation in a southern Brazil Acrisol managed under no-tillage for 17 years. *Soil Tillage Res.* **2005**, *81*, 87–95. [[CrossRef](#)]
69. González-Rosado, M.; Parras-Alcántara, L.; Aguilera-Huertas, J.; Benítez, C.; Lozano-García, B. Effects of land management change on soil aggregates and organic carbon in Mediterranean olive groves. *Catena* **2020**, *195*, 104840. [[CrossRef](#)]
70. Akça, E.; Aydemir, S.; Kadir, S.; Eren, M.; Zucca, C.; Günal, H.; Previtali, F.; Zdruli, P.; Çilek, A.; Budak, M. (Eds.) *The Soils of Turkey, World Soils Book Series*; Springer International Publishing: Cham, Switzerland, 2018; pp. 139–167. [[CrossRef](#)]
71. Polat, O.; Kapur, S. Soil quality parameters in Arenosols under Stone Pine (*Pinus Pinea* L.). Plantations in the Turan Emeksiz sand dune Area. Çukurova University Institute of Science. *J. Sci. Eng.* **2010**, *3*, 167–174. (in Turkish).
72. Yaktı, S.; Akça, E.; Kapur, S. Management of coastal sand dunes by an agroecosystem approach. In *Workshop Proceedings of Determining an Income-Product Generating Approach for Soil Conservation Management*; Zdruli, P., Liuzzi, G.T., Hikmet, E.F., Eds.; CIHEAM: Marrakech, Morocco, 2004; pp. 227–234.
73. Akça, E.; Bal, Y.; Kapur, S.; Eswaran, H. Human-induced late Holocene degradation of the Asi (Orontes) delta, Samandag, Southern Turkey. In *Land Degradation and Rehabilitation: Dryland Ecosystems. In Proceedings of the Fourth International Conference on Land Degradation, Cartagena, Murcia, Spain, 12–17 September 2004*; Catena Verlag: Schweizerbart, Germany, 2004; pp. 17–26, ISBN 3-923381-54.
74. Maia, S.M.F.; Ogle, S.M.; Cerri, C.C.; Cerri, C.E.P. Changes in soil organic carbon storage under different agricultural management systems in the Southwest Amazon Region of Brazil. *Soil Tillage Res.* **2010**, *106*, 177–184. [[CrossRef](#)]
75. Gargouri, K.; Bentaher, H.; Rhouma, A. A novel method to assess drought stress of olive tree. *Agron. Sustain. Dev.* **2012**, *32*, 735–745. [[CrossRef](#)]
76. Chaker, R.; Gargouri, K.; Ben Mbarek, H.; Baraket, F.; Maktouf, S.; Soua, N.; Khelifi, M.; Bouzid, J. Effect of biochemical fraction of exogenous organic matter on CO<sub>2</sub> emission from arid soil: Original Research Article: Effect of biochemical fraction of exogenous organic matter on CO<sub>2</sub> emission from arid soil. *Greenh. Gas Sci. Technol.* **2018**, *8*, 721–733. [[CrossRef](#)]
77. Arshad, M.; Khedher, K.M.; Ayed, H.; Mouldi, A.; Moghanm, F.S.; El Ouni, M.H.; Benkahla, N.; Laatar, E.; Bilal, M.; Abdel Zaher, M. Effects of land use and cultivation histories on the distribution of soil organic carbon stocks in the area of the Northern Nile Delta in Egypt. *Carbon Manag.* **2020**, *11*, 341–354. [[CrossRef](#)]
78. Ogle, S.M.; Breidt, F.J.; Paustian, K. Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry* **2005**, *72*, 87–121. [[CrossRef](#)]
79. Corbeels, M.; Scopel, E.; Cardoso, A.; Bernoux, M.; Douzet, J.M.; Neto, M.S. Soil carbon storage potential of direct seeding mulch-based cropping systems in the Cerrados of Brazil: Soil c storage under dam in the cerrados. *Glob. Chang. Biol.* **2006**, *12*, 1773–1787. [[CrossRef](#)]
80. Don, A.; Schumacher, J.; Freibauer, A. Impact of tropical land-use change on soil organic carbon stocks—a meta-analysis: Soil organic carbon and land-use change. *Glob. Chang. Biol.* **2011**, *17*, 1658–1670. [[CrossRef](#)]