

**Deriving Predictive Relationships for Carotenoid Content at the Canopy Level in a
Conifer Forest using Hyperspectral Imagery and PROSPECT-DART Simulations**

Rocío Hernández-Clemente^{1,*}, Rafael M. Navarro-Cerrillo¹, Pablo J. Zarco-Tejada²

¹E.T.S.I.A.M.-Departamento de Ingeniería Forestal, Universidad de Córdoba, Campus
de Rabanales, 14071 Córdoba, Spain

²Instituto de Agricultura Sostenible (IAS), Consejo Superior de Investigaciones
Científicas (CSIC), Alameda del Obispo, s/n, 14004 Córdoba, Spain

Corresponding author:

Rocío Hernández-Clemente
1E.T.S.I.A.M.-Departamento de Ingeniería Forestal, Universidad de Córdoba,
Campus de Rabanales, 14071 Córdoba, Spain
14004 - Córdoba
Spain
Tel: +(34) 957 218776
Fax: +(34) 957 211 010
e-mail: rociohc@uco.es
www.treesat.com

Abstract

Recent studies have demonstrated that the R_{570}/R_{515} index is highly sensitive to $Cx+c$ content in conifer forest canopies and is scarcely influenced by structural effects. Yet, validated methods for the prediction of leaf carotenoid content relationships in forest canopies are still needed to date. This paper focuses on the simultaneous retrieval of chlorophyll ($Ca+b$) and carotenoid ($Cx+c$) pigments, which are critical bio-indicators of plant physiological status. Radiative transfer theory and modeling assumptions were applied at both laboratory and field scales to develop methods for their concurrent estimation using high-resolution hyperspectral imagery. The proposed methodology was validated based on the biochemical pigment quantification of $Ca+b$ and $Cx+c$ content. Canopy modeling methods based on infinite reflectance formulations and more complex 3-D approaches based on the Discrete Anisotropic Radiative Transfer (DART) model were evaluated in relation to the PROSPECT-5 leaf model for the scaling-up procedure. Simpler modeling methods yielded comparable results to more complex 3-D approximations due to the high spatial resolution images acquired, which enabled targeting pure crowns and reducing the effects of canopy architecture. The scaling-up methods based on the PROSPECT-5+DART model yielded root mean square errors (RMSE) and a relative RMSE of $1.48 \mu\text{g}/\text{cm}^{-2}$ (17.45%) and $5.03 \mu\text{g}/\text{cm}^{-2}$ (13.25%) for $Cx+c$ and $Ca+b$, respectively, while the simpler approach based on the PROSPECT-5+Hapke infinite reflectance model yielded $1.37 \mu\text{g}/\text{cm}^{-2}$ (17.46%) and $4.71 \mu\text{g}/\text{cm}^{-2}$ (14.07%) for $Cx+c$ and $Ca+b$, respectively. These predictive algorithms proved to be useful to estimate $Ca+b$ and $Cx+c$ from high-resolution hyperspectral imagery, providing a methodology for the monitoring of these photosynthetic pigments in conifer forest canopies.

Index Terms: chlorophyll, carotenoids, hyperspectral, airborne, R_{515}/R_{570} , R_{750}/R_{710} , TCARI/OSAVI, conifers, forest, scaling-up

I. INTRODUCTION

Photosynthetic pigments have been identified as important bio-indicators of plant physiological state, mainly due to their roles in photosynthesis [1]-[2]. Early diagnosis of forest decline processes has previously been made using photosynthetic pigment content as a stress indicator [3]-[4]. The main short-term physiological response of trees undergoing decline is a general reduction in photosynthesis due to a large decrease in the levels of chlorophylls and carotenoids [5]. Several authors e.g. [6], [7] have demonstrated that decline in chlorophyll content is associated with an increase in the values of the Ca/Cb ratio. By contrast, the $Ca+b/Cx+c$ ratio shows a decreasing trend under these conditions. Several authors [8], [9] have demonstrated that the increase of needle chlorosis and the decrease rate of carotenoids concentration are associated with decline processes of conifer forest.

The above-mentioned decrease in chlorophyll and carotenoid content in leaves leads to a lower overall absorption in the 430-700 nm region due to a reduction in the proportion of light-absorbing pigments. For this reason, several narrow-band optical indices have been reported to be sensitive to chlorophyll content at the leaf level (see [10], for a review of vegetation indices) and the canopy level [11]. Carotenoid-sensitive indices have been also analyzed, but they have received less attention. Although some authors have proposed indices sensitive to carotenoid pigment at the leaf level [12]-[13], a recent study has explored their sensitivity when applied at the crown and canopy levels [14]. Despite the extensive work conducted on broadleaf species, strategies to link

biochemical and optical properties in coniferous species have been limited by difficulties in measuring coniferous species. The main spectral region proposed for the retrieval of chlorophyll content is located in the red edge, where band ratios are highly sensitive to pigment content [15]. The red edge region shows the maximum slope in vegetation reflectance spectra, between 680 and 750 nm. This is because chlorophyll absorption is dramatically reduced from the red to the near-infrared region. In forest canopies, one of the most sensitive red edge formulations was found by [11], who demonstrated that the R_{750}/R_{710} red edge index was less sensitive than other chlorophyll indices to forest shadows, minimizing their disturbing effects on the retrieval of chlorophyll content. Later, the same red edge index formulation was also validated by other authors (e.g. [16]-[17]). Combined indices sensitive to $Ca+b$ content have recently been developed. An example is the Transformed Chlorophyll Absorption in Reflectance Index, TCARI [18] combined with the Optimized Soil-Adjusted Vegetation Index, OSAVI [19], which minimizes soil effects and is robust for low leaf area index (LAI) levels in crops [18]-[20]-[21]-[22]. Nevertheless, these potentially valuable indices that are suitable for field crops [18], olive orchards [21] and vineyards [20]-[22] have not been validated for forest canopies, whose architecture varies substantially from that of agricultural areas.

Several methods have been proposed to retrieve biophysical parameters from hyperspectral reflectance in forest canopies, including scaling-up and model inversion methods that couple leaf and canopy transfer models [11]- [23]- [16]. Radiative transfer models have been widely used to simulate the bidirectional reflectance distribution of vegetation canopies considering different canopy complexities [24]. There are also simpler simulation approximations of the canopy such as infinite reflectance models

[25]- [2] and [26]. Such models express an optically thick medium in terms of single reflectance and transmittance and consider different assumptions related to the scattering between layered leaves. A previous study reported by [11] suggested that these models are able to estimate pigment content of broadleaf species in closed forests when pure crowns can be targeted using very high resolution imagery. Even under these conditions, the applicability of these simpler methods to the retrieval of pigment content in heterogeneous conifer forest canopies requires further research.

In an effort to model complex forest structures, 3-D models such as SPRINT [27]-[28], FLIGHT [27] or DART [28] have been developed [24]. The application of such models to simulate forest scenes, and in particular the assessment of canopy-level indices for chlorophyll content estimation in conifers using PROSPECT+SPRINT [23], yielded a root square error of $8.1 \mu\text{g}/\text{cm}^2$ when targeting sunlit crown pixels. A more complex approach such as the Discrete Anisotropic Radiative Transfer model (DART) [29] was developed to simulate heterogeneous coniferous forests. As an example, the coupled PROSPECT+DART model was used by [30] to estimate chlorophyll content using an artificial neural network (ANN), yielding a root square error of $2.95 \mu\text{g}/\text{cm}^2$ when using the Area under curve Normalized to maximal Chlorophyll absorption Band depth between 650-720 nm method (ANCB₆₅₀₋₇₂₀).

Despite the efforts made to assess several $Ca+b$ -related vegetation indices, further research is needed to study $Cx+c$ -sensitive indices both at the leaf and canopy levels. Moreover, both pigments exhibit overlapping absorption wavebands, increasing the difficulty of devising a non-destructive method for independently estimating pigment content in plants. For this purpose, we focused on the assessment of different narrow-band vegetation indices sensitive to $Cx+c$ in previous research [14]. In that study, the

R_{515}/R_{570} index was proposed for forestry sites, as it proved to be significantly related to leaf C_{x+c} concentration both at leaf level ($r^2 > 0.72$; $P < 0.001$) and canopy level ($r^2 > 0.71$; $P < 0.001$); yet, it was not possible to estimate C_{x+c} using predictions or scaling-up relationships that accounted for the structure of the canopy. In forest sites, it is essential to be able to quantify C_{a+b} and C_{x+c} independently with sensitive narrow-band indices both at the leaf and canopy levels and develop predictive relationships. Therefore, the main focus of this study was to develop a methodology specifically validated for heterogeneous forest canopies to simultaneously quantify C_{a+b} and C_{x+c} using scaling-up approaches rather than just assess the relationship between indices and carotenoid concentration. The study involved comparing modeling methods using the leaf PROSPECT model coupled with three infinite reflectance models (the simple approach) and a complex 3-D DART model for the retrieval of chlorophyll and carotenoid content at the crown level.

II. METHODS

A. Field experiments and data collection

Field data collection was conducted in August 2008 in Sierra de Filabres (Almeria province, southeastern Spain) ($37^\circ 13' 27''$ N, $2^\circ 32' 54''$ W) (Fig. 1 a), the driest region in Western Europe. The elevation of the study area ranged from 1540 to 2000 m.a.s.l. and annual rainfall ranged between 300 and 400 mm. The annual average temperature was 11°C , reaching a maximum of 32°C in summer and a minimum of -8°C in winter. The vegetation consisted of a 40-year-old pine afforestation of *Pinus nigra* Arnold (Black pine) and *Pinus sylvestris* L. (Scotch pine) (Table 1). Forest decline processes have been previously reported in Sierra de Filabres [31]. The field sampling campaigns

were conducted concurrently with airborne overflights during the last week of July 2008. Needles were collected from the top of the crown by selecting branches of illuminated areas from a total of 21 trees. Needles were frozen in liquid nitrogen in the field and later stored at -80°C prior to determination of chlorophyll *a* and *b* and carotenoids. Mean crown pigment was calculated from a total of 10 young needles (one year-old needles) collected from the top of the crown.

Needle pigment concentration was determined as reported by [32]. Pigment extracts were obtained from a mixed sample of 5 cm of needle material, using 1 linear cm per needle. The area was calculated by assuming that the needle was a half cylinder and the diameter was the measured width of each needle. Needle diameter was measured with a digital caliper precision instrument. Five additional needle samples were used to take structural measurements (thickness and width) to determine water content and dry mass. The needles were ground in a mortar on ice with liquid nitrogen and diluted in acetone up to 5 ml (in the presence of Na ascorbate). After that, the extracts were filtered through a 0.45- μ m filter to separate the pigment extracts from the Na ascorbate. Spectrophotometric determinations were then conducted on the same extracts.

B. Airborne campaigns

The airborne campaign was conducted by the Spanish National Institute of Aerospace Technology (INTA) with an Airborne Hyperspectral Scanner (AHS) (Sensytech Inc., currently Argon St. Inc., Ann Arbor, MI, USA) in the last week of July 2008. Airborne data were obtained at 12:00 GMT, acquiring 2 m spatial resolution imagery in 38 bands in the 0.43-12.5 μ m spectral range. The Field of View (FOV) and Instantaneous Field of View (IFOV) of the AHS sensor were 90° and 2.5 mrad respectively, and study plots

were imaged in the central region of the scene to avoid edge effects. At-sensor radiance processing and atmospheric correction were performed at the INTA facilities. Atmospheric correction was conducted with ATCOR4 based on the MODTRAN radiative transfer model, using aerosol optical depth measured at 550 nm with a Micro-Tops II sun photometer (Solar Light, Philadelphia, PA, USA) at the time of the flight.

The hyperspectral imagery acquired enabled pure crown identification of Scotch pine and Black pine (Fig. 1 b). Single-crown reflectance spectra were extracted using an object-based crown identification algorithm applied to the hyperspectral imagery (Fig. 2 a, b, c). This made it possible to extract the mean hyperspectral reflectance for sunlit and shaded crown components (Fig. 2 d). Vegetation indices (Table 2) were calculated from the crown spectra for the retrieval of chlorophyll $Ca+b$ and carotenoid content $Cx+c$. The $Ca+b$ -related vegetation indices were selected based on previous studies in forest canopies, using simple ratio indices located in the red edge region [16]- [11] and including recent $Ca+b$ -related indices validated for crop canopies such as the TCARI/OSAVI [18]. The $Cx+c$ -sensitive vegetation index used was based on previous work where the R_{515}/R_{570} index was proposed by Hernández-Clemente *et al.* [14] and proved to be significantly related to $Cx+c$ concentration both at the leaf ($r^2>0.72$, $p<0.001$) and canopy ($r^2>0.71$, $p<0.001$) levels.

C. Modeling the retrieval of chlorophyll and carotenoid content

The PROSPECT-5 model was used to simulate needle reflectance and transmittance for varying chlorophyll $Ca+b$ (10-60 $\mu\text{g}/\text{cm}^2$), carotenoid $Cx+c$ (2-16 $\mu\text{g}/\text{cm}^2$) and leaf water (0.01-0.03 cm) content and fixed values of dry matter content and leaf internal structure. Previous studies [16]-[23] have demonstrated the feasibility of the

PROSPECT model to simulate reflectance and transmittance of needles compared to other specific conifer models such as LIBERTY [33]. Table 3 summarizes the nominal values used for leaf level modeling. Fig. 3 shows the simulated spectral response obtained for C_{x+c} variation, between 2 and 16 $\mu\text{g}/\text{cm}^{-2}$, and C_{a+b} values of 15 and 45 $\mu\text{g}/\text{cm}^{-2}$ (Fig. 3 a, b). Both figures show the specific absorption produced by the carotenoid in the spectral region from 500 to 550 nm. The effect of C_{a+b} content variation on the R_{515}/R_{570} carotenoid index calculated at the leaf level is also shown in detail (Fig. 3 c, d). These figures show that R_{515}/R_{570} is affected by chlorophyll concentration variation.

The simulated leaf reflectance and transmittance spectra were used as input for the canopy radiative transfer simulations proposed in this study. The scaling-up analysis was conducted by comparing the accuracy obtained with different canopy reflectance approximations to evaluate the suitability of each model for simultaneous chlorophyll and carotenoid pigment content retrieval. To achieve this purpose, scaling-up methods were first evaluated based on simpler formulae based on infinite reflectance R_{∞} simulations, which did not account for the canopy structure and the viewing geometry effects. The use of this type of models is justified in the high-resolution imagery chosen, which enabled the identification of pure crown vegetation pixels.

After that, a more complex 3-D canopy reflectance radiative transfer model was analyzed and compared to the previous modeling results. This approach has successfully been tested in several studies [23]- [30] and aims at estimating canopy biochemistry with simpler methods, minimizing structural effects through robust indices using models with a reduced number of inputs. The R_{∞} formulations used in this study

were based on i) [2] ($R_{\infty 1}$), ii) [26] ($R_{\infty 2}$) and iii) [25] ($R_{\infty 3}$). In addition, the 3-dimensional Discrete Anisotropic Radiative Transfer (DART) model [29] [34] was used to simulate 3-D vegetation canopies. The DART model has been used to simulate forest canopy reflectance in Norway spruce (*Picea abies*) for the retrieval of chlorophyll content [30] and Scotch pine canopies for analyzing $Cx+c$ -related vegetation indices [14]. In this study, the DART model was used to simulate canopy scenes modeling pure crown reflectance for Scotch and Black pine canopies. These high-resolution 3-D simulations enabled the extraction of simulated sunlit crown reflectance from single trees. The 3-D forest scenes were built using various structural inputs within the range of variation (Table 3) observed from field measurements conducted at the study sites. Fig. 4 shows a sample area acquired with airborne hyperspectral imagery (Fig. 4 a) and the corresponding DART scene simulation (Fig. 4 b). The airborne-AHS and DART-simulated spectra for sunlit and shaded crown reflectance are also shown (Fig. 4 c).

We used a dataset of spectral canopy reflectance generated from the DART radiative transfer model and the three infinite reflectance models proposed to obtain predictive relationships between $Ca+b$ and $Cx+c$ and the TCARI/OSAVI, R_{750}/R_{710} and R_{515}/R_{570} indices. The database of synthetic spectra was generated using 250 simulations with random inputs for $Ca+b$, $Cx+c$, N, LAI, soil reflectance and C_m (nominal values are shown in Table 3). In total, 150 synthetic spectra were used for modeling, conducted with infinite reflectance formulae and through DART canopy simulations. We used 100 additional synthetic spectra to evaluate the Root Mean Square Error (RMSE) and the relative RMSE (R-RMSE) obtained for each scaling-up approach. The regression analyses between $Ca+b$, $Cx+c$ and the sensitive indices were based on quadratic

models. The R_{515}/R_{570} , R_{750}/R_{710} and TCARI/OSAVI indices were included as shown in Equations (1) and (2).

$$Cx+c=f(R_{515}/R_{570}) \quad (1a)$$

$$Cx+c=f(R_{515}/R_{570}, (R_{515}/R_{570})^2, R_{750}/R_{710}) \quad (1b)$$

$$Cx+c=f(R_{515}/R_{570}, (R_{515}/R_{570})^2, TCARI/OSAVI) \quad (1c)$$

$$Ca+b=f(R_{750}/R_{710}) \quad (2a)$$

$$Ca+b=f(R_{750}/R_{710}, (R_{750}/R_{710})^2) \quad (2b)$$

$$Ca+b=f(TCARI/OSAVI, (TCARI/OSAVI)^2) \quad (2c)$$

III. RESULTS

A. Modeling results

The relationships obtained between the simulations conducted with the different approximations through infinite reflectance R_{∞} models and the coupled PROSPECT-5+DART models yielded significant results ($p<0.001$) for $Cx+c$ and $Ca+b$ estimation.

The predictive power of the different canopy approximations applied for the retrieval of pigment content ($Ca+b$, $Cx+c$) yielded coefficients of determination ranging from $r^2=0.6$ to $r^2=0.9$ for both $Cx+c$ (Table 4) and $Ca+b$ (Table 5). $R_{\infty 1}$ and $R_{\infty 2}$ showed slightly better performance than $R_{\infty 3}$ formulations for both $Cx+c$ (Table 4) and $Ca+b$ (Table 5). As regards $Cx+c$ retrieval, the best RMSE values (RMSE=1.35) were obtained by using scaling-up methods based on the model regression (Equation 1b) (Table 4). We observed a consistent improvement in predictive power when using model Equations 1b and 1c compared to model Equation 1a. The differences between these results are explained because models 1b and 1c considered chlorophyll effects for

carotenoid estimation by using both $Cx+c$ and $Ca+b$ sensitive vegetation indices in scaling-up relationships. As an example of the predictive power of the indices applied, the RMSE obtained with the $R_{\infty 1}$ scaling method was $2.11 \mu\text{g}/\text{cm}^2$ (14.73%) using only the simple R_{515}/R_{570} ratio (model 1a) and $1.35 \mu\text{g}/\text{cm}^2$ (10.71%) using the R_{515}/R_{570} and R_{750}/R_{710} vegetation indices simultaneously (model 1b) (Table 4). For $Ca+b$ retrieval, the best coefficient of determination and RMSE values were also obtained by using the $R_{\infty 1}$, $R_{\infty 2}$ scaling methods and model 2b (Table 5). The simulation demonstrated that the overall RMSE obtained using the R_{750}/R_{710} index was lower than that obtained using the TCARI/OSAVI. When the $R_{\infty 1}$ scaling-up method was used, the RMSE improved from 6.97 to $4.41 (\mu\text{g}/\text{cm}^2)$ (15.22 to 9.58%) (Table 5). Among the different canopy reflectance approximations, the use of quadratic equations based on the R_{750}/R_{710} index (model 2b) successfully improved the RMSE values of the estimated $Ca+b$ content.

Using 3-D model simulations, the best results for the estimation of $Cx+c$ were also obtained using the R_{515}/R_{570} and R_{750}/R_{710} vegetation indices simultaneously (Equation 1b); the best results for the estimation of $Ca+b$ were obtained using the R_{750}/R_{710} vegetation index (Equation 2b). A comparison of results obtained with the infinite reflectance models and 3-D model simulations showed similar values in both cases. For example, when a simpler approach such as the $R_{\infty 1}$ was used, results yielded a RMSE of $1.35 \mu\text{g}/\text{cm}^2$ (10.71%) for $Cx+c$ and $4.41 \mu\text{g}/\text{cm}^2$ (9.58%) for $Ca+b$; when the 3-D DART model was used, the RMSE was $1.7 \mu\text{g}/\text{cm}^2$ for $Cx+c$ (11.31%) and $5.31 \mu\text{g}/\text{cm}^2$ (11.50%) for $Ca+b$.

B. Experimental results

The R_{515}/R_{570} , R_{750}/R_{710} and TCARI/OSAVI indices were assessed as a first approach with the airborne imagery to study their sensitivity to $Ca+b$ and $Cx+c$ measured from the study sites. The relationship between $Cx+c$ content and R_{515}/R_{570} extracted from single crowns identified in the airborne imagery showed significant results ($p<0.01$), yielding $r^2=0.57$ for Scotch and Black pine aggregated samples (Fig. 5). The relationships between $Ca+b$ content and the R_{750}/R_{710} (Fig. 6a) and TCARI/OSAVI indices (Fig. 6b) showed that the coefficient of determination was higher for the R_{750}/R_{710} index ($r^2=0.63$) than for the TCARI/OSAVI ($r^2=0.5$). In fact, the TCARI/OSAVI was more affected by the crown structure than the R_{750}/R_{710} index ($p<0.01$ in both cases).

$Ca+b$ and $Cx+c$ content was then estimated by scaling-up the optical indices using the prediction algorithms developed in the previous section. The models obtained from the simulation analysis (Table 4, 5) were applied using the pure-crown reflectance extracted from the airborne imagery for each study site. Table 6 shows the results obtained for the estimation of both $Ca+b$ and $Cx+c$ pigments at canopy level for Scotch pine stands using the four modeling approximations used and the different indices assessed. According to the results, among all the canopy approximations studied, the estimation of $Cx+c$ yielded a RMSE ranging from 1.35 to 2.12 $\mu\text{g}/\text{cm}^2$ (16.94-21.028%) when using quadratic models based on the R_{515}/R_{570} and R_{750}/R_{710} indices (Table 6). The model performance for $Ca+b$ using the R_{750}/R_{710} index yielded an accuracy ranging between 4.41 and 9.04 $\mu\text{g}/\text{cm}^2$ (13.67- 24.57%). Although the best results were obtained when PROSPECT-5 was coupled to DART (RMSE=4.41 $\mu\text{g}/\text{cm}^2$ (13.67%) for $Ca+b$ and RMSE=1.35 $\mu\text{g}/\text{cm}^2$ (12.52%) for $Cx+c$), results obtained using the simpler $R_{\infty 3}$ approach obtained comparable results (RMSE=5.5 $\mu\text{g}/\text{cm}^2$ for $Ca+b$ (32.48%) and

RMSE=1.45 $\mu\text{g}/\text{cm}^2$ for $Cx+c$ (16.94%). The analysis carried out by introducing the TCARI/OSAVI into the model for both $Cx+c$ and $Ca+b$ pigment retrieval showed significantly worse results, yielding poorer RMSE values for all models. Using the TCARI/OSAVI index, the best results obtained for Scotch pine stands through the $R_{\infty 3}$ infinite reflectance simulation yielded values of RMSE=22.76 and 4.78 $\mu\text{g}/\text{cm}^2$ respectively for $Ca+b$ and $Cx+c$ estimations.

Consistently, the results obtained for Black pine stands (Table 7) were similar between infinite reflectance models and DART. For this species, the $R_{\infty 3}$ approach using the R_{515}/R_{570} and R_{750}/R_{710} indices yielded slightly better results, with RMSE=3.75 $\mu\text{g}/\text{cm}^2$ for $Ca+b$ (15.36%) and RMSE=1.17 $\mu\text{g}/\text{cm}^2$ for $Cx+c$ (21.245%). In agreement with the results obtained for Scotch pine, use of such models introducing the TCARI/OSAVI index yielded poorer RMSE values for Black pine, ranging from 18.93 to 26.94 $\mu\text{g}/\text{cm}^2$ for $Ca+b$, and from 1.45 to 2.74 $\mu\text{g}/\text{cm}^2$ for $Cx+c$ (Table 7).

Among the data analyzed for both species, Fig. 7 and Fig 8 show the relationships between image-estimated and field-measured $Cx+c$ and $Ca+b$ pigment retrieval, respectively. In both cases, the lowest RMSE was obtained with the $R_{\infty 3}$ model. For the estimation of $Cx+c$, the RMSE obtained with the $R_{\infty 3}$ model was 1.37 $\mu\text{g}/\text{cm}^2$ (17.45%), while DART, $R_{\infty 1}$ and $R_{\infty 2}$ yielded higher RMSE values (RMSE=1.48 $\mu\text{g}/\text{cm}^2$ (17.45%), 2.00 $\mu\text{g}/\text{cm}^2$ (18.30%) and RMSE=1.97 $\mu\text{g}/\text{cm}^2$ (18.20%), respectively). For the estimation of $Ca+b$, the RMSE obtained with the $R_{\infty 3}$ model was 4.71 $\mu\text{g}/\text{cm}^2$ (14.07%) while DART, $R_{\infty 1}$ and $R_{\infty 2}$ yielded higher RMSE values (RMSE=5.03 $\mu\text{g}/\text{cm}^2$ (13.25%), 11.95 $\mu\text{g}/\text{cm}^2$ (28.89%) and RMSE =11.85 $\mu\text{g}/\text{cm}^2$ (28.90%), respectively).

The higher performance of $R_{\infty 3}$ in forest canopies agrees with the results obtained by [11], who demonstrated that model inversion using $R_{\infty 3}$ +PROSPECT showed lower RMSE compared to SAILH+PROSPECT using R_{750}/R_{710} in the merit function if high-resolution imagery was available. It is worth noting the high performance obtained in forest canopies with simpler infinite reflectance models based on Hapke's approach compared to the results obtained with more complex and computationally expensive approaches. These results may be explained by the fact that pure crown reflectance was extracted based on the high resolution of the images; therefore, soil and shadow effects had a lower influence on target reflectance. The prediction models assessed applied at image level using object-based methods enabled the mapping of $Cx+c$ and $Ca+b$ with both the R_{515}/R_{570} and R_{750}/R_{710} vegetation indices acquired from the AHS hyperspectral imagery (Fig. 9). Mapping results made it possible to estimate both pigments at the crown level for the entire scene showing the spatial variability of carotenoid content, with areas with high $Cx+c$ content and low $Cx+c$ content (Fig. 9).

IV. DISCUSSION

This study provides insight into the quantification of chlorophyll and carotenoid content simultaneously on conifer forest canopies using scaling-up approaches and narrow-band indices. The major advantage of the methodology used is avoiding the problems of overlapping between wavebands that absorb both pigments (450-550 nm range) [35]. The improved retrieval of carotenoids and chlorophyll contents was obtained combining two narrow-band indices sensitive to both pigments respectively: the R_{515}/R_{570} index proposed by [14] for Scotch pine sites and the red edge index (R_{750}/R_{710}). In the latter case, the optical indices evaluated and related to $Ca+b$ have been analyzed by other authors [16]- [17]- [11]- [23] for conifer forests. This study demonstrated that better

results were obtained when considering the R_{750}/R_{710} index for $Ca+b$. This result may be explained by the higher effect that tree shadow pixels produce on the TCARI/OSAVI index. This agrees with [21], who demonstrated that the TCARI/OSAVI index is highly affected by direct soil background and shadow components. These results differ from the relationship obtained between the TCARI/OSAVI and canopy $Ca+b$ content for herbaceous and open-tree orchard canopies [18]- [21]- [22]. By contrast, successful results were obtained with the R_{750}/R_{710} index by [16], who analyzed $Ca+b$ content using hyperspectral observations in Jack Pine stands, and by [11], who studied the same pigment in Sugar Maple stands.

The estimation of pigment composition in forest canopies using scaling-up methods implies the analysis of the potential confounding effects of open canopy structures on narrow-band vegetation indices. For this reason, the use of complex 3-D radiative transfer modeling was developed to simulate heterogeneous forest canopies [34]. However, the results presented in this study show that simpler canopy approximations can be also applied targeting pure crown spectral information. These results agree with those obtained by [36], who demonstrated the feasibility of scaling-up methods based on the infinite reflectance approach for crops. In fact, the accuracy of scaling-up methods applied to achieve $Ca+b$ and $Cx+c$ estimations based on infinite reflectance approaches is remarkable. Despite our efforts analyzing complex 3-D radiative transfer models to include the structural variations found in conifer forests, simpler approaches ($R_{\infty 3}$) yielded similar results. This was due to the fact that very high resolution was used to extract pure crown reflectance from high spatial resolution hyperspectral imagery, removing mixed pixels, shadows and background effects. It is important to highlight that the forest canopy analyzed in this study had a relatively low heterogeneity, as it was the product of a systematic afforestation with very low species mix. Therefore, the

results obtained with the infinite reflectance approach and the DART simulation analysis might vary in other types of canopies, where the identification of pure crowns is more complicated due to the vegetation mixture or structural heterogeneity of the canopy. This implies that further studies should be carried out in more complex forest canopies comparing different simulation reflectance approaches as a function of varying pixel resolutions.

So far, the methodology proposed in this study for the simultaneous quantification of both pigments ($Ca+b$ and $Cx+c$) is especially relevant when vegetation stress is addressed. This enables the assessment of the spatial and temporal variations of those pigments previously demonstrated to be related to forest decline processes [8], [9]. Moreover, an accurate spatial quantification of pigment content estimations is presented in this work using high resolution hyperspectral image information. Therefore, the methodology proposed in this study for the simultaneous estimation of both pigments is likely to be useful to analyse spatial and temporal declining physiological states.

V. CONCLUSION

This paper builds on previous work focused on indices related to carotenoid content estimation. In this study, scaling-up and predictive relationships were developed using radiative transfer models. Our results demonstrated the feasibility of simultaneously estimating needle $Ca+b$ and $Cx+c$ with scaling-up methods using hyperspectral airborne reflectance data acquired from conifer canopies. These results were obtained based on the scaling up of pure crown vegetation indices related to chlorophyll (R_{750}/R_{710}) and carotenoids (R_{515}/R_{570}) by using modeling simulations conducted with infinite reflectance models based on Hapke's model coupled with PROSPECT-5. The modeling and experimental results were obtained using the R_{515}/R_{570} and R_{750}/R_{710}

vegetation indices for the simultaneous estimation of C_{x+c} and C_{a+b} . The better performance of the R_{750}/R_{710} index as compared to the TCARI/OSAVI index for chlorophyll content retrieval in forest canopies was probably due to the greater shadow effects of a heterogeneous architecture such as a forest stand on the TCARI/OSAVI. The accuracy obtained by applying scaling-up methods that used simpler approaches such as the infinite reflectance formulation proposed by Hapke was comparable to more complex canopy reflectance approximations such as the DART model. This was due to the very high resolution imagery used, which enabled pure crown identification. Under these conditions, the modeling analysis and experimental measurements were conducted for the simultaneous estimation of pigment content at the crown level, yielding mean errors of $1.37 \mu\text{g}/\text{cm}^2$ (17.45%) for C_{x+c} retrieval, and $4.71 \mu\text{g}/\text{cm}^2$ for C_{a+b} retrieval (14.07%). In addition, the promising results obtained with 3-D model simulation demonstrate the suitability of this methodology for more complex forest canopies. More complex models such as the 3-D DART canopy reflectance model estimated C_{a+b} and C_{x+c} content slightly more accurately than simpler infinite reflectance models; they showed mean absolute errors of $1.48 \mu\text{g}/\text{cm}^2$ (17.45%) for C_{x+c} retrieval and $5.03 \mu\text{g}/\text{cm}^2$ for C_{a+b} retrieval (13.25%), but required a considerably larger number of inputs. The generation of biochemical maps at the crown level could play a critical role in the early detection of forest decline processes, enabling the application of such models in precision forestry.

ACKNOWLEDGEMENTS

This project was funded through DIVERBOS (CGL2011-30285-C02-02). We also acknowledge the financial support from University of Córdoba-Campus de Excelencia CEIA3.). Support from the Spanish Ministry of Science and Education (MEC) to the

AGL2009-13105 project is gratefully acknowledged. We are also grateful to J. E. Granados and the members of CAHA (German-Spanish Astronomical Center at Calar Alto), R. Sanchez (ERSAF), D.Ariza (IDAF), E. Gil-Pelegrín and J. J. Peguero-Pina (CITA, Zaragoza) for providing technical support during the fieldwork campaigns. Finally, we greatly appreciate the support of J. P. Gastellu-Etchegorry with the DART 3-D radiative transfer model.

REFERENCES

- [1] G.A. Carter, and R.L. Miller, “Early detection of plant stress by digital imaging within narrow stress-sensitive wavebands,” *Remote Sens. Environ.*, vol. 50, pp. 295-302. Dec, 1994.
- [2] O. Lillesaeter, “Spectral reflectance of partly transmitting leaves: Laboratory measurements and mathematical modeling,” *Remote Sens. Environ*, vol. 12, pp. 247-254. Jul. 1982
- [3] F. Batič, P. Kalan, H. Kraigher, H. Šircelj, P. Simončič, N. Vidergar-Gorjup, and B. Turk, “Bioindication of different stresses in forest decline studies in Slovenia,” *Water, Air and Soil Pollution*, vol. 116, pp. 377-382, Feb. 1999.
- [4] R. Hernández-Clemente, R.M. Navarro-Cerrillo, L. Suárez, F. Morales, and P.J. Zarco-Tejada, “Assessing structural effects on PRI for stress detection in conifer forests,” *Remote Sens. Environ*, vol. 115, pp. 2360-2375, Sep. 2011.
- [5] M. Lippert, K.Steiner, H.D. Payer, S. Simons, C. Langebartels, and H. Sandermann, “Assessing the impact of ozone on photosynthesis of European beech (*Fagus sylvatica* L.) in environmental chambers,” *Trees*, vol. 10, pp. 268–275, Jun. 1996.

- [6] T. Hoshizaki, B. N. Rock, and S. K. S. Wong, "Pigment analysis and spectral assessment of spruce trees undergoing forest decline in the NE United States and Germany". *GeoJournal*, vol. 17, pp. 173-178, Sep.1988.
- [7] R. Matyssek, M.S. Günthardt-Goerg, T. Keller and C. Scheidegger, "Impairment of gas exchange and structure in birch leaves (*Betula pendula*) caused by low ozone concentrations," *Trees*, vol.5, pp. 5-13, Nov. 1991.
- [8] T. Batič, Z. Užarevic, L. Grgic, J. Roša, and Z. Popovic, "Chlorophylls and carotenoids in needles of damage fir (*Abies alba* Mill.) from Risnjak National Park in Croatia. *Acta biologica cracoviensia*," vol. 45, pp. 87–92, Jun. 2003.
- [9] R. Oren, K.S. N. Werk, Buchmann, and R. Zimmermann, "Chlorophyll–nutrient relationships identify nutritionally caused decline in Piceaabies stands," *Canadian Journal of Forest Research*, vol. 23, pp. 1187-1195, Jun. 1993.
- [10] R. Main, M. A. Cho, R. Mathieu, M. O’Kennedy, M. A. Ramoelo, and S. Koch, "An investigation into robust spectral indices for leaf chlorophyll estimation," *ISPRS Journal of Photogrammetry and Remote Sensing* , vol. 66, pp. 751-761, Nov. 2011.
- [11] P. J. Zarco-Tejada, J. R. Miller, G. H. Mohammed, T. L. Noland, and P. H. Sampson, "Scaling-up and Model Inversion methods with narrow-band Optical Indices for Chlorophyll Content Estimation in closed Forest Canopies with Hyperspectral Data". *IEEE Transactions on Geoscience and Remote Sensing*, vol. 39, pp. 1491-1507, Jul, 2001.
- [12] A. A. Gitelson, Y. Zur, O. B Chivkunova, and M. N. Merzlyak, "Assessing carotenoid content in plant leaves with reflectance spectroscopy," *Journal of Photochemistry and Photobiology B-Biology*, vol. 75, pp. 272-281, Dec. 2002.

- [13] J. Peñuelas, I. Filella, and Gamon, J. A., “Assessment of photosynthetic radiation use efficiency with spectral reflectance,” *New Phytologist*, vol. 131, pp. 291-296, Nov. 1995.
- [14] R. Hernández-Clemente, R. M. Navarro-Cerrillo, and P. J. Zarco-Tejada, “Carotenoid content estimation in an heterogeneous conifer forest using narrow-band indices and PROSPECT + DART simulations,” *Remote Sens. Environ*, vol. 127, pp. 298-315, Sep. 2012.
- [15] D. N. H., Holder, M Dockray, and J. Barber, “The red edge of plant leaf reflectance,” *International Journal of Remote Sensing*, vol. 4, pp. 273-278, 1983.
- [16] I. Moorthy, J. R. Miller, and T. L. Noland, “Estimating chlorophyll concentration in conifer needles with hyperspectral data: An assessment at the needle and canopy level,” *Remote Sens. Environ*, vol. 112, pp. 2824-2838, Jun. 2008.
- [17] Q. Zhang, X. Xiao, B. Braswell, E. Linder, F. Baret, and B. Moore, “Estimating light absorption by chlorophyll, leaf and canopy in a deciduous broadleaf forest using MODIS data and a radiative transfer model,” *Remote Sens. Environ*, vol. 99, pp. 357-371, Nov. 2005.
- [18] D. Haboudane, J. R. Miller, N. Tremblay, P. J. Zarco-Tejada, and L. Dextraze, (2002), “Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture,” *Remote Sens. Environ*, vol. 81, pp. 416-426, Aug. 2002.
- [19] G. Rondeaux, M. Steven, and F. Baret, “Optimization of soil-adjusted vegetation indices,” *Remote Sens. Environ*, vol. 55, pp. 95-107, Feb. 1996.

- [20] F. Meggio, P. J. Zarco-Tejada, L.C. Núñez, G. Sepulcre-Cantó, M. R. Gonzalez, and P. Martin, "Grape quality assessment in vineyards affected by iron deficiency chlorosis using narrow-band physiological remote sensing indices," *Remote Sens. Environ*, vol. 114, pp. 1968-198, Sep. 2010.
- [21] P. J. Zarco-Tejada, J. R. Miller, A. Morales, A. Berjón, and J. Agüera, "Hyperspectral Indices and Model Simulation for Chlorophyll Estimation in Open-Canopy Tree Crops," *Remote Sens. Environ*, vol. 90, pp. 463-476, Apr. 2004.
- [22] P. J. Zarco-Tejada, A. Berjón, R. López-Lozano, J. R. Miller, P. Martín, V. Cachorro, M. R. González, and A. Frutos, "Assessing Vineyard Condition with Hyperspectral Indices: Leaf and Canopy Reflectance Simulation in a Row-Structured Discontinuous Canopy," *Remote Sens. Environ*, vol. 99, pp. 271-287, Nov. 2005.
- [23] P. J. Zarco-Tejada, J. R. Miller, J. Harron, B. Hu, T. L. Noland, N., Goel, G. H. Mohammed, and P. H. Sampson (2004), "Needle chlorophyll content estimation through model inversion using hyperspectral data from boreal conifer forest canopies," *Remote Sens. Environ*, vol. 89, pp. 189-199, Jan. 2004.
- [24] B. Pinty, J. L. Widlowski, M. Taberner, N. Gobron, M. M. Verstraete, M. Disney, P. Lewis, F. Gascon, J. P. Gastellu, L. Jiang, X. Li, L. Su, S. Tang, H. Wang, J. Wang, G. Yan, H. Zang, A. Kuusk, T. Nilson, W. Ni-Meister, P. North, W. Qin, R. Thompson, and W. Verhoef (2004), "Radiation Transfer Model Intercomparison (RAMI) exercise: Results from the second phase," *Journal of Geophysical Research D: Atmospheres*, vol. 109, pp. 1-19, Mar. 2004.
- [25] B. Hapke, *Theory of Reflectance and Emittance Spectroscopy*. Cambridge, U.K. Univ. Press, 1993.

- [26] N. Yamada and S Fujimura, "Non-destructive measurement of chlorophyll pigment content in plant leaves from three-color reflectance and transmittance". *Appl. Opt.*, vol. 30, pp. 3964-3973, Sep. 1991.
- [27] P. R. J. North, (1996), "Three dimensional forest light interaction model using a Monte Carlo method," *IEEE Transactions on Geosciences and Remote Sensing*, vol. 34, pp. 946–956, Jul. 1996.
- [28] N. S. Goel, and R. L. Thompson, "A snapshot of canopy reflectance models and a universal model for the radiation regime," *Remote Sensing Reviews*, vol. 18, pp. 197–22, 2000.
- [29] J. P. Gastellu-Etchegorry, E. Martin, and F. Gascon, (2004), "DART: a 3-D model for simulating satellite images and studying surface radiation budget," *International Journal of Remote Sensing*, vol. 25, pp. 73-96.
- [30] Z. Malenovsky, L. Homolova,, P. Cudlin,, R. Zurita Milla,, M.E. Schaepman, J.G.P.W. Clevers, E. Martin, and J.P. Gastellu-Etchegory, "Physically-based retrievals of Norway spruce canopy variables from very high spatial resolution hyperspectral data". In *Geoscience and Remote Sensing Symposium (IGARSS)*, Vol. 1 (pp. 4057-4060). Barcelona (Sp): IEEE, July. 2007.
- [31] R. Sánchez-Salguero, R. Navarro-Cerrillo, T. W. Swetnam, and M. A. Zavala. "Is drought the main decline factor at the rear edge of Europe?: the case of southern Iberian pine plantations?," *For Ecol Manage.*, vol. 271, pp.158-169, May. 2012.
- [32] A. Abadía, and J. Abadía, "Iron and plant pigments," in *Iron Chelation in Plants and Soil Microorganisms*, L. L. Barton and B. C. Hemming ed., Academic: San Diego, 1993, pp. 327-344.

- [33] T. P. Dawson, P. J. Curran, and S. E. Plummer (1998), “LIBERTY- modelling the effects of leaf biochemistry on reflectance spectra,” *Remote Sens. Environ*, vol. 65, pp. 50- 60, Jul. 1998.
- [34] J. P. Gastellu-Etchegorry, V. Demarez, V. Pinel, and Zagolski, F. (1996), “Modeling radiative transfer in heterogeneous 3-D vegetation canopies,” *Remote Sens. Environ*, vol. 58, pp. 13-156, Nov. 1996.
- [35] Y. Zur, A.A. Gitelson, O.B. Chivkunova, and M.N. Merzlyak, “The spectral contribution of carotenoids to light absorption and reflectance in green leaves,” in Proceedings of the 2nd International Conference Geospatial Information in Agriculture and Forestry, Buena Vista, FL, January 10-12, 2000, v. 2.
- [36], P. J. Zarco-Tejada, M. L. Guillén-Climent, R. Hernández-Clemente, A. Catalina, M. R. González, and P. Martín, “Estimating carotenoid content in vineyards using high resolution hyperspectral imagery acquired from an unmanned aerial vehicle (UAV),” *Agr.Forest Meteorol.*, vol. 171-172, pp. 281-294, Apr. 2013.



Rocío Hernández Clemente, obtained an Agricultural Engineering Degree at the Faculty of Agricultural Engineering and Forestry, University of Córdoba (UCO) (Spain), an International PhD degree accredited by the Department of Forestry (University of Cordoba)- Her specific lines of research are focused on the analysis of high-resolution hyperspectral data to estimate leaf biochemical and canopy biophysical variables through leaf and canopy radiative transfer modeling. She is currently at the Treesat laboratory (UCO) (Spain), where she has a contract as scientific researcher of the FP7 Thermolidar project

based on the application of advanced technologies and new methods to improve estimates of structural information and physiological variables in forestry based on the fusion of airborne LiDAR with thermal data sets.



Pablo J. Zarco-Tejada holds a degree in agricultural engineering from the School of Agricultural Engineering and Forestry, University of Córdoba, Spain, an MSc degree in remote sensing from the University of Dundee, Scotland, UK, and a PhD degree in earth and space science from York University, Toronto, ON, Canada. He has been a contract faculty member for remote sensing at the University of California, Davis. He works on hyperspectral data to estimate biochemical and canopy biophysical variables through leaf and canopy modeling and the effects of chlorophyll fluorescence. He is currently with the Institute for Sustainable Agriculture, Spanish Council for Scientific Research, Córdoba, Spain.



Rafael Maria Navarro Cerrillo, professor at the university of Cordoba and Head of the ERSAF research group. He has more than 25 years' experience on forestry science. He has 20 years of working experience in silviculture, restauration and remote sensing. He has participated in different international projects in Europe, Latin America and Asia.